

Exercise not shown to reduce women's risk of developing multiple sclerosis

September 28 2016

A large, new study shows no evidence that exercise may reduce a woman's risk of developing multiple sclerosis (MS). The research is published in the September 28, 2016, online issue of *Neurology*, the medical journal of the American Academy of Neurology. Previous small studies had shown conflicting results.

"We wanted to find out if [exercise](#) lowered the risk of developing MS in women," said study author Kassandra Munger, ScD, of the Harvard T.H. Chan School of Public Health in Boston, Mass. "Our study did not provide evidence to support it."

Researchers evaluated data on more than 193,000 women who participated in the Nurses' Health Study and Nurses' Health Study II and were followed for up to 20 years. The women completed regular questionnaires about their physical activity and also about their activity as teens and young adults. During the study, 341 women developed MS.

Researchers calculated the total hours of [physical activity](#) per week, took into account the type of exercise for each woman and adjusted for age, ethnicity, smoking, supplemental vitamin D, place of residence at age 15 and [body mass index](#) at age 18.

"Overall, there was no consistent association of exercise at any age and MS," Munger said. "Exercise has been shown to be beneficial to people with the disease, but it seems unlikely that exercise protects against the risk of developing MS."

Provided by American Academy of Neurology

Citation: Exercise not shown to reduce women's risk of developing multiple sclerosis (2016, September 28) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-09-shown-women-multiple-sclerosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.