


Sitagliptin not linked to higher fracture risk in diabetes

September 14 2016



(HealthDay)—Patients with diabetes often have fractures, but sitagliptin



is not associated with increased fracture risk, according to a study published online Sept. 8 in *Diabetes, Obesity and Metabolism*.

Robert G. Josse, M.B.B.S., from the University of Toronto, and colleagues examined fracture incidence in the Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS). Data were included from 14,671 TECOS patients who were randomized to sitagliptin (7,332 patients) or placebo (7,339 patients).

The researchers found that 2.6 percent of patients had a fracture during 43,222 person-years' follow-up, of which 146 were major osteoporotic fractures. There was an independent association between fracture risk and older age, female sex, white race, and lower diastolic blood pressure (all P fracture risk was seen in association with insulin therapy (hazard ratio, 1.4; P = 0.035), and metformin was linked to decreased risk (hazard ratio, 0.76; P = 0.035).

"Fractures were common among TECOS patients with diabetes, but not related to sitagliptin therapy," the authors write. "Insulin and metformin treatment were associated with increased and decreased fracture risks, respectively."

Several authors disclosed financial ties to pharmaceutical companies, including Merck, which manufactures sitagliptin and funded the study.

More information: <u>Abstract</u>

Full Text (subscription or payment may be required)

Copyright © 2016 HealthDay. All rights reserved.

Citation: Sitagliptin not linked to higher fracture risk in diabetes (2016, September 14) retrieved 26 April 2024 from



https://medicalxpress.com/news/2016-09-sitagliptin-linked-higher-fracture-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.