

# Specific trauma experiences contribute to women's alcohol use, differs by race

September 22 2016

---

Trauma exposure has consistently been reported as a risk factor for alcohol use and related problems. Further, racial differences in alcohol use, alcohol use disorder (AUD), and trauma exposure between European American (EA) and African American (AA) women have been reported previously. This study sought to identify racial differences in alcohol involvement, and to examine the risk conferred by specific trauma exposures and PTSD for different stages of alcohol involvement in EA and AA women.

Researchers examined data from the Missouri Adolescent Female Twins Study: the mean age of the 3,787 women at time of interview was 24.5 years; 85.4 percent were EA, 14.6 percent AA. Trauma exposures were defined as sexual abuse (SA), physical abuse (PA), witnessing another person being killed or injured, experiencing an accident, or experiencing a disaster. Trauma exposure was examined as a predictor of alcohol initiation, transition to the first AUD symptom, and transition to an AUD diagnosis – while also considering other substance involvement, parental characteristics, and commonly co-occurring psychiatric disorders.

Results showed that trauma experiences were important contributors to all stages of alcohol involvement in EA women only, with different trauma types conferring risk for each stage of alcohol involvement. For example, in EA women SA was associated with alcohol initiation prior to the age of 14; PA predicted the transition from initiation to the first AUD symptom; and PA, witnessing injury or death, and SA predicted

the transition to an AUD diagnosis. There were no such findings in AA women. Further, PTSD was not revealed as a significant predictor of AUD in either EA or AA women. The findings suggest that trauma, independent of PTSD, directly contributes to alcohol involvement. Further, they highlight the importance of considering [racial differences](#) when looking at linkages between traumatic experiences and alcohol involvement.

**More information:** Kimberly B. Werner et al. Association of Specific Traumatic Experiences With Alcohol Initiation and Transitions to Problem Use in European American and African American Women, *Alcoholism: Clinical and Experimental Research* (2016). [DOI: 10.1111/acer.13220](#)

Provided by Research Society on Alcoholism

Citation: Specific trauma experiences contribute to women's alcohol use, differs by race (2016, September 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-09-specific-trauma-contribute-women-alcohol.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--