

Activity trackers are ineffective at sustaining weight loss, study finds

September 20 2016, by Anthony Moore



The new FitBit Blaze

Wearable devices that monitor physical activity are not reliable tools for weight loss, says a new study from the University of Pittsburgh School of Education's Department of Health and Physical Activity. The study specifically investigated whether regular use of commercially available activity trackers is effective for producing and sustaining weight loss.

At the conclusion of a 24-month trial, researchers observed that usage of a wearable device in combination with a behavioral [weight loss](#) program resulted in less weight loss when compared to those receiving only the behavioral [weight loss program](#). In fact, participants without physical activity trackers showed nearly twice the weight loss benefits at the end of the 24 months. Participants who utilized wearable devices reported an average weight loss of 7.7 pounds, while those who partook only in health counseling reported an average loss of 13 pounds.

Through these observations, researchers concluded that devices that monitor and provide feedback on physical activity do not offer an advantage over standard weight loss approaches that include behavioral counseling on physical activity and diet. Thus, while these devices allow for ease of tracking of physical activity along with feedback and encouragement, they may not enhance adherence to the tenets of a healthy lifestyle, which is the most important aspect of any weight loss regiment.

"While usage of wearable devices is currently a popular method to track physical activity—steps taken per day or calories burned during a workout—our findings show that adding them to behavioral counseling

r weight loss that includes physical activity and reduced calorie intake does not improve weight loss or physical activity engagement. Therefore, within this context, these devices should not be relied upon as tools for weight management in place of effective behavioral counseling for physical activity and diet," said John Jakicic, the study's lead researcher and chair of Pitt's Department of Health and Physical Activity.

The study "Effect of Wearable Technology Combined with a Lifestyle Intervention on Long-Term Weight Loss: the IDEA Randomized Clinical Trial," was released today in the latest edition of *JAMA: The Journal of the American Medical Association*.

The study followed 470 individuals between the ages of 18 and 35 with a body mass index between 25 and 39 at the start of the trial.

Approximately 77 percent of participants were women and 29 percent were from minority communities.

All participants were placed on low-calorie diets, prescribed increases in physical activity, and received group-counseling sessions on health and nutrition. They participated in weekly health-counseling sessions for the initial six months and less frequent counseling for the last 18 months. Weight was assessed at six-month intervals throughout the 24-month trial.

At the first six-month mark, participants were divided into two subgroups: one that continued health-counseling sessions on a monthly basis and another that received a [wearable device](#) to monitor diet and physical activity. The multisensor device used within the study was to be worn on the upper arm and provided feedback on energy expenditure and physical activity. A website was designed to monitor results.

Over the course of the subsequent 18 months, both groups showed significant improvements in body composition, fitness, physical activity, and diet, with no significant difference between groups. However, those who received health counseling throughout the study lost nearly twice as much weight as those who used wearable devices for three-quarters of it.

"The findings of our study are important because effective long-term treatments are needed to address America's obesity epidemic," said Jakicic. "We've found that questions remain regarding the effectiveness of wearable devices and how to best use them to modify [physical activity](#) and diet behaviors in adults seeking weight loss."

More information: Effect of Wearable Technology Combined With a Lifestyle Intervention on Long-term Weight Loss, *JAMA*.

2016;316(11):1161-1171. [DOI: 10.1001/jama.2016.12858](https://doi.org/10.1001/jama.2016.12858) ,
[jama.jamanetwork.com/article.a ... px?articleid=2553448](https://jamanetwork.com/article.a...px?articleid=2553448)

Provided by University of Pittsburgh

Citation: Activity trackers are ineffective at sustaining weight loss, study finds (2016, September 20) retrieved 20 March 2024 from <https://medicalxpress.com/news/2016-09-trackers-ineffective-sustaining-weight-loss.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--