

Study: Care providers underestimate pain during pediatric burn dressing change

September 1 2016

Dressing changes have been identified as one of the major contributors to perceived pain during burn care, and so it is important that pain be assessed accurately. But a new study from the Center for Pediatric Trauma Research at Nationwide Children's Hospital suggests that pediatric burn pain assessment can vary not only based on patient pain intensity, but also nurse clinical experience.

A measurement tool called the Face, Legs, Activity, Cry, Consolability Scale (FLACC) is widely used by nurses to rate [pain](#) in pediatric patients. It can be applied to pediatric patients of all ages, including the very young, so it provides a standard assessment for all patients in our pediatric burn center. This study found that nurses tended to underrate both mild and severe procedural pain. Nurses who had less nursing experience demonstrated significantly higher accuracy than those with more experience.

Jonathan Groner MD is the study's senior author, medical director of the Center for Pediatric Trauma Research and affiliate faculty of the Center for Injury Research and Policy at Nationwide Children's Hospital stated, "This is concerning because it demonstrates that on a well-established burn unit that specializes in caring for [pediatric patients](#), nurses may too often underrate children's [procedural pain](#) for both mild and severe pain." Dr. Jiabin Shen, the study's lead author also noted concern that "new" burn nurses (less than 5 years nursing experience) appeared to be more accurate in FLACC rating than more experienced [nurses](#).

This study, available today online and in the September issue of *Burns*, is the first study to systematically examine FLACC rating accuracy among care providers. Developed in 1997, the FLACC scale was originally designed to measure post-operative pain in pediatric populations. The FLACC utilizes an easy-to-understand 0-10 metric and exerts a relatively low burden on medical staff. It has previously demonstrated a high level of inter-rater reliability and concurrent validity with other existing scales. But much of the previous research was not in clinical settings.

Provided by Nationwide Children's Hospital

Citation: Study: Care providers underestimate pain during pediatric burn dressing change (2016, September 1) retrieved 6 May 2024 from <https://medicalxpress.com/news/2016-09-underestimate-pain-pediatric.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--