

Study investigates absorption of artificial sweeteners in blood

October 24 2016



Credit: CC0 Public Domain

A recent study by investigators at the National Institute of Diabetes,

Digestive and Kidney Diseases at the National Institutes of Health measured how much artificial sweetener is absorbed into the blood stream by children and adults after drinking a can of diet soda. Results of this study are published in *Toxicological & Environmental Chemistry*.

The team measured the artificial sweeteners sucralose and acesulfame-potassium, which are found in a wide range of packaged foods and beverages. These artificial sweeteners, also including saccharin and aspartame, have received a lot of attention lately because it has been found that they are not inert chemicals with a sweet taste, but active substances that can affect the metabolism.

Despite their approval as food additives following the submission of detailed safety data to the United States Food and Drug Administration (FDA), concerns about their safety and especially about their long-term [health](#) effects remain. Artificial sweetener use is increasing worldwide because it is universally accepted that high sugar consumption promotes a variety of health problems, including obesity and [diabetes](#). The food industry responds to the consumer demand, and increasingly replaces sugar with artificial sweeteners in order to provide tasty goods with lower sugar content. Most consumers expect that weight loss will result from switching to artificial sweeteners (because they contain no or fewer calories), but paradoxically the opposite may happen.

Given this background, the authors performed a study to extend previous investigations into plasma concentrations of sucralose and acesulfame-potassium. Artificial sweetener concentrations were measured among [adults](#) following ingestion of various doses of sucralose with or without acesulfame-potassium, both in diet soda and mixed in seltzer or plain water. Results obtained in adults were then compared with measurements obtained in children.

The study comprised 22 adults aged 18-45 and 11 children aged 6-12

with no known medical conditions and who were not using any medications, enrolled in a randomized same-subject crossover study. The protocol was approved by the Institutional Review Board of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).

The results of the study demonstrate that, compared to adults, children had double the concentrations of plasma sucralose after ingestion of a single twelve ounce can [diet soda](#). The same research team previously found that these artificial sweeteners were also present in breast milk when mothers ingested foods, drinks, medicines or other products that contained artificial sweeteners. Since infants have less ability to clear substances from their [blood stream](#) via the kidneys (lower glomerular filtration rate up to age 2 years), the authors speculate that the infants' [artificial sweetener](#) blood levels may be proportionately even higher.

These results are important because early life exposure to artificial sweeteners may influence a child's future taste preferences, diet and metabolic fate. Since it is known that [children](#) generally prefer more sweetness than adults, they are especially vulnerable to the intense sweetness provided by artificial sweeteners.

The findings of this study also highlight the fact that some people absorb relatively small amounts of artificial sweeteners and some exorbitantly high amounts. Overall, this study will help others with the design and interpretation of crucial future research to better understand what artificial sweeteners do to our health.

More information: Allison C. Sylvetsky et al. Plasma concentrations of sucralose in children and adults, *Toxicological & Environmental Chemistry* (2016). [DOI: 10.1080/02772248.2016.1234754](https://doi.org/10.1080/02772248.2016.1234754)

Provided by Taylor & Francis

Citation: Study investigates absorption of artificial sweeteners in blood (2016, October 24)
retrieved 3 May 2024 from

<https://medicalxpress.com/news/2016-10-absorption-artificial-sweeteners-blood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.