

More Americans undergo procedures involving anesthesia outside of O.R.

October 22 2016

More than one-third of Americans who undergo procedures involving anesthesia now have them outside of the operating room (O.R.), an increase of 27 percent in five years, according to an analysis of a large registry being presented at the Anesthesiology 2016 annual meeting. Researchers say these patients tend to be older and in poorer health, having been urged by physicians to have less-risky minimally invasive procedures and tests that don't need to be performed in an O.R.

"The growth reflects a shift toward safer, less-invasive procedures and tests that still require anesthesia," said Richard Dutton, M.D., a senior author of the study and chief quality officer for U.S. Anesthesia Partners, Dallas. "Whether inside or outside an O.R., it's important that anesthesia for patients with severe health conditions is provided by physician anesthesiologists who have the extensive training and skill to deliver safe care."

The research is based on an analysis of the Anesthesia Quality Institute's National Anesthesia Clinical Outcomes Registry (NACOR) of anesthesia cases in all settings, from hospitals to freestanding clinics. Researchers identified more than 5.9 million non-O.R. anesthesia cases and 12.4 million O.R. cases from 2010 through 2014 and determined the percentage of cases performed outside the O.R. increased from 28 percent to 36 percent. Non-O.R. patients were an average of 3.5 years older than O.R. patients. Colonoscopy (a screening test for colon cancer) was the most common non-O.R. procedure involving anesthesia.

Procedures that involve large incisions typically take place in an O.R., a sterile environment with extensive specialized equipment, and often are performed under [general anesthesia](#), which makes the patient lose consciousness. Conversely, non-O.R. procedures often are performed by inserting a catheter into an artery, through tiny incisions or using a scope through the mouth or rectum and may be less risky than surgery, particularly for older patients who have heart disease, stroke, severe liver disease or other life-threatening conditions.

Anesthesia provided during non-O.R. procedures is more likely to involve intravenous (IV) sedation that may range from minimal (making the patient drowsy but able to talk) to deep (the patient likely won't remember the procedure) or regional [anesthesia](#) that numbs part of the body. These procedures can be performed in radiological suites or other procedure rooms that don't feature the same extensive equipment as the O.R.

Provided by American Society of Anesthesiologists

Citation: More Americans undergo procedures involving anesthesia outside of O.R. (2016, October 22) retrieved 2 May 2024 from <https://medicalxpress.com/news/2016-10-americans-procedures-involving-anesthesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
