

# Risk of frailty associated with changes in oral function

October 3 2016

---



(HealthDay)—Aging is tied to changes in oral function, according to a

study published online Sept. 22 in the *Journal of the American Geriatrics Society*.

Yutaka Watanabe, D.D.S., Ph.D., from the Tokyo Metropolitan Institute of Gerontology, and colleagues conducted detailed physical testing to classify elderly adults ( $\geq 65$  years) by frailty (mobility, strength, endurance, physical activity, and nutrition): robust, prefrail, or frail. Additionally, the numbers of present teeth and functional teeth were counted, plus measurements of occlusal force, masseter muscle thickness, and oral diadochokinesis (ODK) rate.

The researchers found that the frail group had significantly fewer teeth present (women aged  $\geq 70$ ), lower occlusal force (women aged  $\geq 70$ ; men aged  $\geq 80$ ), lower masseter muscle thickness, and lower ODK rate, compared to the robust group. Multivariate analysis indicated that frailty was also significantly associated with age, Geriatric Depression Scale score, skeletal muscle mass index, Mini-Mental State Examination score, hypertension, diabetes mellitus, albumin and triglyceride levels, and oral function.

"The risk of frailty was associated with lower occlusal force, masseter muscle thickness, and ODK rate," the authors write.

**More information:** [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Risk of frailty associated with changes in oral function (2016, October 3) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-10-frailty-oral-function.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.