


Improved leg ulcer healing with hair follicle punch graft

October 25 2016



(HealthDay)—For patients with venous leg ulcers, ulcer healing is

significantly increased using hair follicle punch scalp grafts, according to a study published in the November issue of the *Journal of the American Academy of Dermatology*.

María-Luisa Martínez, M.D., from the General Hospital of Villarrobledo in Albacete, Spain, and colleagues conducted a randomized trial with intraindividual comparison of [hair follicle](#) scalp grafts and nonhairy skin grafts, which were transplanted in parallel into two halves of the same ulcer in 12 patients.

The researchers found that at 18 weeks after the intervention, ulcer healing, as measured by the average percentage reduction, was increased significantly in the hair follicle group, with a reduction in the ulcer area of 75.15 percent compared with 33.07 percent in the control group (nonhairy grafts) ($P = 0.002$).

"Autologous transplantation of terminal hair follicles by scalp punch grafts induces better healing than punch grafts harvested from nonhairy areas," the authors write. "Hair punch grafting is a minimally [invasive surgical procedure](#) that appears to be effective as a therapeutic tool for chronic [venous leg ulcers](#)."

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Improved leg ulcer healing with hair follicle punch graft (2016, October 25) retrieved 8 May 2024 from <https://medicalxpress.com/news/2016-10-leg-ulcer-hair-follicle-graft.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.