

Study suggests patch is safe, convenient mode of treatment for peanut allergy

October 26 2016



Viaskin Peanut patch used in the NIH-sponsored clinical trial. The Viaskin Peanut patch has not been approved by the FDA. Credit: DBV Technologies, Copyright 2016

A wearable patch that delivers small amounts of peanut protein through the skin shows promise for treating children and young adults with



peanut allergy, with greater benefits for younger children, according to one-year results from an ongoing clinical trial. The treatment, called epicutaneous immunotherapy or EPIT, was safe and well-tolerated, and nearly all participants used the skin patch daily as directed.

The ongoing trial is sponsored by the National Institute of Allergy and Infectious Diseases (NIAID), part of the National Institutes of Health, and conducted by the NIAID-funded <u>Consortium of Food Allergy</u> <u>Research</u> (CoFAR), which is led by Hugh Sampson, M.D., of Icahn School of Medicine at Mount Sinai in New York. Stacie Jones, M.D., of the University of Arkansas for Medical Sciences and Arkansas Children's Hospital, chairs the study. One-year outcomes are published online on October 26 in the *Journal of Allergy and Clinical Immunology*.

"To avoid potentially life-threatening allergic reactions, people with peanut allergy must be vigilant about the foods they eat and the environments they enter, which can be very stressful," said NIAID Director Anthony S. Fauci, M.D. "One goal of experimental approaches such as epicutaneous immunotherapy is to reduce this burden by training the immune system to tolerate enough peanut to protect against accidental ingestion or exposure."

CoFAR researchers at five study sites randomly assigned 74 peanutallergic volunteers aged 4 to 25 years to treatment with either a highdose (250 micrograms peanut protein), low-dose (100 micrograms peanut protein), or placebo patch. The investigators assessed peanut allergy at the beginning of the study with a supervised, oral food challenge with peanut-containing food. The patches were developed and provided by the biopharmaceutical company DBV Technologies under the trade name Viaskin. Each day, study participants applied a new patch to their arm or between their shoulder blades.

After one year, researchers assessed each participant's ability to consume



at least 10 times more <u>peanut protein</u> than he or she was able to consume before starting EPIT. The low-dose and high-dose regimens offered similar benefits, with 46 percent of the low-dose group and 48 percent of the high-dose group achieving treatment success, compared with 12 percent of the placebo group. In addition, the peanut patches induced immune responses similar to those seen with other investigational forms of immunotherapy for food allergy. Investigators observed greater treatment effects among children aged 4 to 11 years, with significantly less effect in participants aged 12 years and older.

"The clinical benefit seen in younger children highlights the promise of this innovative approach to treating peanut allergy," said Daniel Rotrosen, M.D., director of NIAID's Division of Allergy, Immunology and Transplantation (DAIT). "Epicutaneous immunotherapy aims to engage the immune system in the skin to train the body to tolerate small amounts of allergen, whereas other recent advances have relied on an oral route that appears difficult for approximately 10 to 15 percent of children and adults to tolerate."

Nearly all of the <u>study participants</u> followed the EPIT regimen as directed. None reported serious reactions to the patch, although most experienced mild skin reactions, such as itching or rash, at the site of patch application.

"The high adherence to the daily peanut patch regimen suggests that the patch is easy-to-use, convenient and safe," said Marshall Plaut, chief of DAIT's Food Allergy, Atopic Dermatitis and Allergic Mechanisms Section. "The results of this study support further investigation of epicutaneous immunotherapy as a novel approach for peanut allergy treatment."

More information: Stacie M. Jones et al. Epicutaneous immunotherapy for the treatment of peanut allergy in children and young



adults, *Journal of Allergy and Clinical Immunology* (2016). DOI: <u>10.1016/j.jaci.2016.08.017</u>

Provided by National Institutes of Health

Citation: Study suggests patch is safe, convenient mode of treatment for peanut allergy (2016, October 26) retrieved 16 June 2024 from <u>https://medicalxpress.com/news/2016-10-patch-safe-convenient-mode-treatment.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.