

## **Pulmonary embolism may be cause of syncope in some elderly**

October 20 2016

---



(HealthDay)—About one of every six patients hospitalized for a first

episode of syncope has a pulmonary embolism, according to a study published in the Oct. 20 issue of the *New England Journal of Medicine*.

Researchers at 11 hospitals in Italy performed a systematic work-up for [pulmonary embolism](#) in 560 patients admitted for a first-time syncope episode. The patients were 76 years old, on average, and had been admitted from the [emergency department](#) for various reasons: The cause of syncope was not apparent; there was reason to suspect a cardiovascular-related cause; they had other serious [medical conditions](#); or they'd been injured when they fainted.

The team found that 17.3 percent—or roughly one in six—were diagnosed with pulmonary embolism. That included 13 percent of patients who'd had a potential alternative explanation for their syncope, such as a cardiovascular condition.

"In [elderly patients](#) presenting with syncope, the attending physician in medical wards should consider pulmonary embolism as a possible differential diagnosis—particularly when an alternative explanation is not found," study coauthor Sofia Barbar, M.D., a physician at the Civic Hospital of Camposampiero in Padua, Italy, told *HealthDay*.

**More information:** [Abstract](#)  
[Full Text](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Pulmonary embolism may be cause of syncope in some elderly (2016, October 20) retrieved 19 April 2024 from <https://medicalxpress.com/news/2016-10-pulmonary-embolism-syncope-elderly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.