

Team questions safety, efficacy of selenium and colorectal cancer risk

October 14 2016



A UA-led clinical trial found that selenium supplementation did not prevent colon polyps and may increase the risk of type 2 diabetes. Credit: University of Arizona Health Sciences

A 12-year study led by a team of University of Arizona Cancer Center researchers is bringing into question the safety and efficacy of selenium, a popular nutritional supplement touted to combat and reduce the risk of colorectal cancer.

The findings indicate the need for a significant change in practice, given that selenium supplements cannot be recommended for preventing




colorectal cancer.

Selenium has been a popular nutritional supplement for decades, touted for its antioxidant properties and its role in stopping free radicals from damaging cells and DNA. Studies have shown a deficiency of this micronutrient to be associated with cancer risk.

However, a <u>randomized clinical trial</u> involving 1,824 participants from clinical centers in Arizona, Colorado, Texas and New York indicates that selenium supplements failed to prevent the development of colon polyps, but significantly increased the risk of developing type 2 diabetes in older individuals.

"The possibility that selenium supplements may increase the risk of type 2 diabetes has been hinted at before," said Peter Lance, MD, deputy director of the UA Cancer Center and the study's principal investigator. "But this is the first study to have substantiated such a risk in the setting of a prospective, randomized, placebo-controlled trial." Study participants were randomized to take 200 mcg of selenium as selenized yeast or a placebo daily.





Cancer—Histopathologic image of colonic carcinoid. Credit: Wikipedia/CC BY-SA 3.0

The trial also looked at celecoxib, a selective COX-2 inhibitor non-steroidal anti-inflammatory drug (NSAID), versus placebo. While the once-daily 400 mg dose prevented colon polyps in high-risk patients, it increased the risk of hypertension in those with pre-existing cardiovascular risk factors. "Our study has added important refinements to understanding who may benefit most from the colon polyp-preventing effects of celecoxib and how best to avoid cardiovascular toxicity caused by this agent," added Dr. Lance.

Further UA-led studies are under way to define the biological mechanisms underlying the effects, including toxicity, of <u>selenium</u> <u>supplements</u>. A study is planned to target celecoxib to those patients who



will benefit most from the colon polyp-preventing actions of celecoxib while minimizing cardiovascular toxicity.

The findings of the Selenium and Celecoxib Trial recently were published online before print in two articles in the *Journal of the National Cancer Institute* (JNCI): "Selenium Supplementation for Prevention of Colorectal Adenomas and Risk of Associated Type 2 Diabetes" and "Celecoxib for the Prevention of Colorectal Adenomas: Results of a Suspended Randomized Controlled Trial". The articles will appear in print in the December 2016 issue of *JNCI*.

More information: Patricia A. Thompson et al, Selenium Supplementation for Prevention of Colorectal Adenomas and Risk of Associated Type 2 Diabetes, *Journal of the National Cancer Institute* (2016). DOI: 10.1093/jnci/djw152

Patricia A. Thompson et al. Celecoxib for the Prevention of Colorectal Adenomas: Results of a Suspended Randomized Controlled Trial, *Journal of the National Cancer Institute* (2016). DOI: 10.1093/jnci/djw151

Provided by University of Arizona

Citation: Team questions safety, efficacy of selenium and colorectal cancer risk (2016, October 14) retrieved 20 March 2024 from https://medicalxpress.com/news/2016-10-team-safety-efficacy-selenium-colorectal.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.