

White coat hypertension may indicate risk for heart disease in some people

October 31 2016

White coat hypertension, where patients have high blood pressure readings in a medical setting but normal blood pressure outside the doctor's office, is most likely an innocuous condition that is not a predictor of heart disease or stroke—except in a small group of older patients, according to a study published online today in the *Journal of the American College of Cardiology*.

White coat hypertension, identified more than 30 years ago, is a common condition that is thought to be the result of patients feeling stressed in a medical setting or in the presence of clinicians in "white coats," though some research has suggested it is associated with health risks. Recent studies have shown white coat hypertension occurs more frequently in [older patients](#) and the difference between the [blood pressure](#) readings in the two settings becomes larger with age.

To find out if white coat hypertension is an indicator of increased risk for [heart disease](#), a researcher from the University of California, Irvine, in collaboration with the International Database on Ambulatory blood pressure in relation to Cardiovascular Outcomes, looked at 653 patients identified with white coat hypertension from the international hypertension database and compared them to 653 patients of similar age and similar risk for developing heart disease.

The researchers, led by Stanley S. Franklin, M.D., found that during a follow up period of about 10.6 years there was no difference in the number of new heart-related health events between younger subjects,

below age 60, with [normal blood pressure](#) and those of a similar age and risk profile with white coat hypertension.

A difference was seen in older patients. Out of 92 high-risk subjects age 60 and older, there were 18 more new cardiovascular events in the follow up period for those diagnosed with white coat hypertension than the same number of subjects of similar age and otherwise similar risk profile with normal blood pressure.

Researchers said the results support the hypothesis that a small number of patients considered to have white coat hypertension actually have isolated systolic hypertension, a common condition where the top number in the blood pressure reading (systolic blood pressure) is too high, but the bottom number (the diastolic blood pressure) is normal. Isolated [systolic hypertension](#) can be an indicator of risk for future heart disease or stroke.

The authors conclude multiple [blood pressure readings](#), including readings outside a clinical setting, are necessary to accurately identify cardiovascular risk, especially in older high-risk patients.

In an accompanying editorial, Giuseppe Mancina, M.D., and Guido Grassi, M.D., of the University of Milano Bicocca, Milan, Italy, said that the study group may not have been large enough or studied over a long enough time period to identify risk in the younger patients, so conclusions about younger patients may not be accurate, and more research is needed to fully understand the complexities of [white coat hypertension](#).

More information: *Journal of the American College of Cardiology*, DOI: [10.1016/j.jacc.2016.08.035](https://doi.org/10.1016/j.jacc.2016.08.035)

Provided by American College of Cardiology

Citation: White coat hypertension may indicate risk for heart disease in some people (2016, October 31) retrieved 5 May 2024 from <https://medicalxpress.com/news/2016-10-white-coat-hypertension-heart-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.