

Study says CT use OK for some colon cancers, yet challenging for lymph node involvement

November 28 2016

CT has good sensitivity for the detection of colon cancers with tumors that have spread beyond the bowel wall, however, it remains a challenge in detecting nodal involvement, which could have considerable consequences given the increasing interest in neoadjuvant treatment for colon cancer.

This finding was published in the November 2016 issue of the *American Journal of Roentgenology (AJR)* in an article titled, "Diagnostic Accuracy of CT for Local Staging of Colon Cancer: A Systematic Review and Meta-Analysis." The article, which is available on *AJR's* website through open access, indicates that thin-slice CT may be an effective tool in identifying nodal involvement.

"Detecting nodal involvement with CT is unreliable," Elias Nerad, radiologist at Catharina Hospital Eindhoven and researcher at Netherlands Cancer Institute in Amsterdam, The Netherlands, said. "The use of thin slices improved the detection of tumor invasion beyond the bowel wall, as well as the detection of malignant lymph nodes, and is therefore advocated. Also, evidence suggests that CT colonography improves the accuracy of CT in the local staging of [colon tumors](#), which could have a major clinical effect if neoadjuvant chemotherapy is adopted in the [treatment](#) of [colon cancer](#)."

CT colonography is a technique that uses optimal bowel wall distention

and 3D reconstruction of the colon (creating a virtual colonoscopy), which distinguishes it from regular CT of the abdomen. The meta-analysis provided by study coauthor Nerad, et al. was limited to two CT colonography studies because the majority of them identified in the literature search included rectal tumors that were inseparable from the colon tumors and thus were excluded.

"CT Colonography is highly sensitive and cost effective in detection of colorectal carcinoma," Nerad said. "Furthermore, it can also be used as a (local) staging tool, making it a very interesting potential subgroup in our meta-analysis."

Overall, the research was based on the meta-analysis of 13 studies describing the accuracy of CT in the staging of colon carcinomas. In the literature, colon and rectal cancers are combined despite the fact that they differ in terms of anatomy, diagnostic workup, and treatment. This study does make a clear distinction between the two and aims to exclusively determine the diagnostic accuracy of CT for staging colon cancer only and to evaluate whether CT can help in differentiating high- from low-risk colon cancer tumors.

Therapy for rectal cancer is based on staging with MRI or endorectal ultrasound, the study said. For the treatment of MRI-staged high-risk rectal tumors, defined as tumors with involvement of the mesorectal fascia, guidelines recommend neoadjuvant chemoradiotherapy.

In contrast, colon cancer is mainly staged with CT and there are no guidelines for the use of neoadjuvant treatment for colon cancer. However, this will most likely change in the near future, if ongoing studies confirm earlier reports that patients with aggressive colon tumors benefit from a neoadjuvant treatment.

The meta-analysis provided by Nerad et al. specifically focuses on the

[diagnostic accuracy](#) of CT for staging colon cancer and to evaluate whether CT can select patients for neoadjuvant treatment (i.e., aggressive colon tumors).

More information: Elias Nerad et al. Diagnostic Accuracy of CT for Local Staging of Colon Cancer: A Systematic Review and Meta-Analysis, *American Journal of Roentgenology* (2016). [DOI: 10.2214/AJR.15.15785](#)

Provided by American Roentgen Ray Society

Citation: Study says CT use OK for some colon cancers, yet challenging for lymph node involvement (2016, November 28) retrieved 11 May 2024 from <https://medicalxpress.com/news/2016-11-ct-colon-cancers-lymph-node.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.