

Reducing exposure to bisphenol A (BPA) lowers levels of this environmental estrogen in women

November 3 2016



Credit: Mary Ann Liebert, Inc., publishers

Women who avoided foods, cosmetics, and other products packaged in BPA-containing plastic containers for 3 weeks had significant reductions in urinary levels of BPA, a commonly used "endocrine disruptor" associated with negative health effects including weight gain. Over the 3-week study period, the women who participated in an intervention designed to minimize BPA exposure also had significant weight loss, as reported in *Journal of Women's Health*.

The article, entitled "Randomized Intervention Trial to Decrease Bisphenol A Urine Concentrations in Women: Pilot Study", shows that among the participants, the women in the control group who did not take part in the BPA-limiting intervention, had significant increases in both urinary BPA levels and [weight gain](#) after 3 weeks.

Todd Hagobian, PhD and coauthors from California Polytechnic State University (San Luis Obispo, CA), propose future large-scale randomized trials to confirm these findings and to determine the potential positive health effects of reduced exposure to endocrine disrupting chemicals on risk factors for disorders such as type 2 diabetes and cardiovascular disease.

"This study shows that by switching to BPA-free products it is, in fact, possible for women who have been exposed to BPA to reduce their body burden of the compound, as measured by urinary BPA levels," says Susan G. Kornstein, MD, Editor-in-Chief of *Journal of Women's Health*, Executive Director of the Virginia Commonwealth University Institute for Women's Health, Richmond, VA, and President of the Academy of Women's Health. "Although many consumers tend to reject products made of plastics containing BPA, there are unfortunately still many other endocrine disrupting chemicals in our environment."

More information: Todd Hagobian et al, Randomized Intervention Trial to Decrease Bisphenol A Urine Concentrations in Women: Pilot Study, *Journal of Women's Health* (2016). [DOI: 10.1089/jwh.2016.5746](https://doi.org/10.1089/jwh.2016.5746)

Provided by Mary Ann Liebert, Inc

Citation: Reducing exposure to bisphenol A (BPA) lowers levels of this environmental estrogen in women (2016, November 3) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-11-exposure-bisphenol-bpa-lowers-environmental.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.