

High-fiber diet keeps gut microbes from eating colon's lining, protects against infection

November 17 2016



In experiments in germ-free mice that received a transplanted set of human gut microbes, a high-fiber diet was associated with a thick mucus layer in the colon's lining, and resistance to infection by a pathogen. But mice raised with a fiber-free diet had a much thinner, patchy mucus layer, as mucus-eating bacteria in the human microbe mix proliferated. The pathogen was able to get through to the cells of the colon wall. Credit: University of Michigan



It sounds like the plot of a 1950s science fiction movie: normal, helpful bacteria that begin to eat their host from within, because they don't get what they want.

But new research shows that's exactly what happens when microbes inside the digestive system don't get the natural fiber that they rely on for food.

Starved, they begin to munch on the natural layer of mucus that lines the gut, eroding it to the point where dangerous invading bacteria can infect the colon wall.

In a new paper in *Cell*, an international team of researchers show the impact of fiber deprivation on the guts of specially raised <u>mice</u>. The mice were born and raised with no <u>gut microbes</u> of their own, then received a transplant of 14 bacteria that normally grow in the <u>human gut</u>. Scientists know the full genetic signature of each one, making it possible to track their activity over time.

The findings have implications for understanding not only the role of fiber in a normal diet, but also the potential of using fiber to counter the effects of digestive tract disorders.

"The lesson we're learning from studying the interaction of fiber, gut microbes and the intestinal barrier system is that if you don't feed them, they can eat you," says Eric Martens, Ph.D., an associate professor of microbiology at the University of Michigan Medical School who led the research along with his former postdoctoral fellow Mahesh Desai, Ph.D., now at the Luxembourg Institute of Health.

Using U-M's special gnotobiotic, or germ-free, mouse facility, and advanced genetic techniques that allowed them to determine which bacteria were present and active under different conditions, they studied



the impact of diets with different fiber content - and those with no fiber. They also infected some of the mice with a bacterial strain that does to mice what certain strains of Escherichia coli can do to humans - cause gut infections that lead to irritation, inflammation, diarrhea and more.

The result: the <u>mucus layer</u> stayed thick, and the infection didn't take full hold, in mice that received a diet that was about 15 percent fiber from minimally processed grains and plants. But when the researchers substituted a diet with no fiber in it, even for a few days, some of the microbes in their guts began to munch on the mucus.

They also tried a diet that was rich in prebiotic fiber - purified forms of soluble fiber similar to what some processed foods and supplements currently contain. This diet resulted in the same erosion of the mucus layer as observed in the lack of fiber.

The researchers also saw that the mix of bacteria changed depending on what the mice were being fed, even day by day. Some species of bacteria in the transplanted microbiome were more common - meaning they had reproduced more - in low-fiber conditions, others in high-fiber conditions.





A thick mucus layer, generated by the cells of the colon's wall, provides



protection against invading bacteria and other pathogens. This image of a mouse's colon shows the mucus (green) acting as a barrier for the "goblet" cells (blue) that produce it. In this case, the mice were raised in a germ-free environment, then received a transplanted set of human gut microbes. When the mice were fed a high-fiber diet, the mucus layer was thick and microbes that can digest fiber flourished. But on a fiber-free diet, the mucus layer became much thinner after microbes capable of digesting mucus became more dominant. This left the goblet cells open to infection by a pathogen. Credit: University of Michigan

And the four <u>bacteria strains</u> that flourished most in low-fiber and nofiber conditions were the only ones that make enzymes that are capable of breaking down the long molecules called glycoproteins that make up the mucus layer.

In addition to looking at the of bacteria based on genetic information, the researchers could see which fiber-digesting enzymes the bacteria were making. They detected more than 1,600 different enzymes capable of degrading carbohydrates - similar to the complexity in the normal human gut.

Just like the mix of bacteria, the mix of enzymes changed depending on what the mice were being fed, with even occasional fiber deprivation leading to more production of mucus-degrading enzymes.

Images of the mucus layer, and the "goblet" cells of the colon wall that produce the mucus constantly, showed the layer was thinner the less fiber the mice received. While mucus is constantly being produced and degraded in a normal gut, the change in bacteria activity under the lowestfiber conditions meant that the pace of eating was faster than the pace of production - almost like an overzealous harvesting of trees outpacing the planting of new ones.



When the researchers infected the mice with *Citrobacter rodentium* - the *E. coli*-like bacteria - they observed that these dangerous bacteria flourished more in the guts of mice fed a fiber-free diet. Many of those mice began to show signs of illness and lost weight.

When the scientists looked at samples of their gut tissue, they saw not only a much thinner or even patchy mucus later - they also saw inflammation across a wide area. Mice that had received a fiber-rich diet before being infected also had some inflammation but across a much smaller area.

Martens notes that in addition to the gnotobiotic facility, the research was possible because of the microbe DNA and RNA sequencing capability built up through the Medical School's Host Microbiome Initiative, as well as the computing capability to plow through all the sequence data.

"Having all the resources here was the key to making this work, and the fact that it was all across the street from our lab allowed us to pin it all together," he says. He also notes the role of U-M colleagues led by Gabriel Nunez and Nobuhiko Kamada in providing the *C. rodentium* pathogen model, and of French collaborators from the Aix-Marseille Université in studying the enzymes in the mouse gut.

Going forward, Martens and Desai intend to look at the impact of different prebiotic fiber mixes, and of diets with more intermittent natural fiber content over a longer period. They also want to look for biomarkers that could tell them about the status of the mucus layer in human guts - such as the abundance of mucus-digesting <u>bacteria</u> strains, and the effect of low fiber on chronic disease such as inflammatory bowel disease.

"While this work was in mice, the take-home message from this work



for humans amplifies everything that doctors and nutritionists have been telling us for decades: Eat a lot of fiber from diverse natural sources," says Martens. "Your diet directly influences your microbiota, and from there it may influence the status of your gut's mucus layer and tendency toward disease. But it's an open question of whether we can cure our cultural lack of fiber with something more purified and easy to ingest than a lot of broccoli."

More information: Cell, DOI: 10.1016/j.cell.2016.10.043

Provided by University of Michigan Health System

Citation: High-fiber diet keeps gut microbes from eating colon's lining, protects against infection (2016, November 17) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2016-11-high-fiber-diet-gut-microbes-colon.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.