

Incidence of sebaceous carcinoma up from 2000 to 2012

November 29 2016



(HealthDay)—From 2000 to 2012 there was an increase in the incidence

of sebaceous carcinoma (SC), according to a study published in the December issue of the *Journal of the American Academy of Dermatology*.

Raghav Tripathi, from Case Western Reserve University in Cleveland, and colleagues analyzed trends in SC in the United States from 2000 to 2012. Data were obtained from 18 Surveillance, Epidemiology, and End Results Program registries and were used to calculate cause of death, relative frequencies/incidences, five-/10-year survival, hazard ratios, and [incidence rates](#) for SC.

The researchers found that overall incidence was 0.32 and 0.16 per 100,000 person-years for males and females, respectively. There was a significant increase in incidence, mainly due to an increase among men. Incidence was almost three times higher among whites than non-whites. Significantly higher all-cause [mortality](#) was seen for male sex, black race, and extraocular anatomic location. There was a significant decrease in overall case-specific mortality for SC.

"The overall incidence of SC is increasing significantly," the authors write. "Male sex, black race, and extraocular occurrences are associated with significantly greater mortality."

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Incidence of sebaceous carcinoma up from 2000 to 2012 (2016, November 29) retrieved 17 May 2024 from <https://medicalxpress.com/news/2016-11-incidence-sebaceous-carcinoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.