

Lack of brain shrinkage may help predict who develops dementia with Lewy bodies

November 2 2016

Dementia with Lewy bodies is a progressive disease that causes hallucinations, decline in mental abilities, rigid muscles, slow movement and tremors. With symptoms similar to Alzheimer's disease and Parkinson's disease, a correct diagnosis can be difficult.

A new study published today in the online issue of *Neurology*, a medical journal of the American Academy of Neurology, shows that a lack of shrinkage in the area of the brain called the hippocampus may be a sign that people with thinking and memory problems may develop dementia with Lewy bodies rather than Alzheimer's disease. Atrophy of the hippocampus, the area of the brain responsible for thinking and memory, is an early sign of Alzheimer's disease.

"Identifying people with mild cognitive impairment at risk for dementia with Lewy bodies is critical for early interventions," says the study's lead author Kejal Kantarci, M.D., a Mayo Clinic radiologist. "Early diagnosis helps target appropriate treatments, including what medications not to give. For example, as many as 50 percent of people with Lewy body disease have severe reactions to antipsychotic drugs."

Lewy bodies are protein deposits that develop in nerve cells in regions of the brain involved in thinking, memory and movement.

In the study, 160 people with [mild cognitive impairment](#) had brain MRI scans to measure hippocampus size. They also had yearly tests for an average of two years. During that time, 61 people, or 38 percent,

developed Alzheimer's disease, and 20 people, or 13 percent, progressed to probable dementia with Lewy bodies. Because Lewy body disease can be diagnosed only by an autopsy after death, it is called probable dementia with Lewy bodies. The researchers note that their results should be confirmed with studies that use autopsies for final diagnoses.

The people who had no shrinkage in the hippocampus were 5.8 times more likely to develop probable dementia with Lewy bodies than those who had hippocampal atrophy. Seventeen of 20, or 85 percent, of people who developed dementia with Lewy bodies had a normal hippocampus volume; whereas, 37 of the 61, or 61 percent, of people who developed Alzheimer's disease had hippocampus atrophy.

The relationship of hippocampus volume and disease was stronger among people without memory issues. Dementia with Lewy bodies does not always affect memory. Affected thinking skills usually include attention, problem-solving and interpreting visual information.

Provided by Mayo Clinic

Citation: Lack of brain shrinkage may help predict who develops dementia with Lewy bodies (2016, November 2) retrieved 10 April 2024 from <https://medicalxpress.com/news/2016-11-lack-brain-shrinkage-dementia-lewy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--