

Do nights, weekends affect survival after pediatric cardiac arrest in hospital?

November 7 2016

For hospitalized children, the rate of surviving to discharge was lower for those who had cardiac arrest with cardiopulmonary resuscitation (CPR) at night compared with during the daytime and evening, according to an article published online by *JAMA Pediatrics*.

Nearly 6,000 children in the United States receive CPR in the [hospital](#) each year and many of these children do not survive to be discharged from the hospital. Some studies of adults have suggested patients have worse outcomes when they have cardiac arrest at night.

Farhan Bhanji, M.D., M.Sc.(Ed.), F.R.C.P.C., of McGill University, Montreal, Canada, and coauthors used the American Heart Association's Get With the Guidelines Resuscitation registry, a large multicenter registry of in-hospital cardiac arrests, to examine survival rates for hospitalized children who had cardiac arrest by the time of day and day of the week.

The study included 12,404 hospitalized children (more than half were male) who received CPR for at least two minutes. Of the children, 8,568 required CPR during daytime or evening hours and 3,836 needed CPR at night. The 354 hospitals who contributed data during the study period had a median size of 333 beds.

Of the 12,404 children, 8,731 (70.4 percent) experienced a return of circulation that lasted more than 20 minutes, 7,248 (58.4 percent) survived for 24 hours and 4,488 (36.2 percent) survived to hospital

discharge, according to the results.

After adjusting for potential mitigating factors, the rate of survival to hospital discharge was about 12 percent lower during nights than during days and evenings but was not different between weekends and weekdays, the authors report.

"Although the absolute rate of survival to hospital discharge was lower on weekends than weekdays, this difference did not reach statistical significance when adjusted for confounding factors," the authors write. The study also acknowledges several limitations, including an inability to identify underlying causes for the differences in survival.

The authors call lower survival rates at nighttime "an important, yet underrecognized public health concern."

"This is especially pertinent because suboptimal resuscitative efforts are a potentially preventable harm. Assuming an annual CPR rate of 6,000 events per year, we found that simply improving overall survival (currently at 36.2%) to match the weekday daytime epoch survival (41.1%) would result in almost 300 additional children's lives saved per year in the United States. These findings may have important implications for hospital staffing, training, and resource allocation," according to the paper.

The study concludes: "Discrepancy between daytime and nighttime outcomes represents an important patient safety concern that warrants further investigation."

More information: *JAMA Pediatr.* Published online November 7, 2016. [DOI: 10.1001/jamapediatrics.2016.2535](https://doi.org/10.1001/jamapediatrics.2016.2535)

Provided by The JAMA Network Journals

Citation: Do nights, weekends affect survival after pediatric cardiac arrest in hospital? (2016, November 7) retrieved 4 May 2024 from <https://medicalxpress.com/news/2016-11-nights-weekends-affect-survival-pediatric.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.