

Poor sensitivity for HPV 16/18 in minor abnormal cytology

November 17 2016



(HealthDay)—Testing for human papillomavirus (HPV) 16/18 has poor

sensitivity for triaging women with minor abnormal cytology, according to a meta-analysis published online Nov. 15 in the *Annals of Internal Medicine*.

Marc Arbyn, M.D., from the Scientific Institute of Public Health in Brussels, and colleagues conducted a systematic literature review to examine the accuracy of genotyping for HPV types 16 and 18. Data were included for 24 moderate-to-good quality studies, involving 8,587 women with atypical squamous cells of undetermined significance (ASC-US) and 5,284 with low-grade squamous intraepithelial lesions (LSIL) who were triaged with tests for high-risk HPV (hrHPV) and HPV 16/18 to detect [cervical intraepithelial neoplasia](#) (CIN).

The researchers found that for women with either ASC-US or LSIL the pooled sensitivity of HPV 16/18 genotyping for CIN3+ was about 70 percent. With a threshold of grade

Citation: Poor sensitivity for HPV 16/18 in minor abnormal cytology (2016, November 17) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2016-11-poor-sensitivity-hpv-minor-abnormal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--