

Anesthesiologist reaction time increased after night call

December 12 2016



(HealthDay)—For anesthesiologists, reaction time increases after a night

call, with greater increases seen in correlation with reliance on avoidance as a coping mechanism, according to a study published online Nov. 30 in *Pediatric Anesthesia*.

Haleh Saadat, M.D., from the Nationwide Children's Hospital in Columbus, and colleagues examined the impact of [sleep deprivation](#) on reaction time in 23 [anesthesiologists](#) after an overnight call or regular working hours. Reaction time was assessed using the psychomotor vigilance test on a regular non-call day and after a 17-hour in-house call. Two anesthesiologists ended their participation for personal reasons.

The researchers found that, compared with a regular day, post-call psychomotor vigilance test mean reaction time decreased by an average of 31.2 ms. Coping Strategy Indicator scale scores indicated [reliance](#) on specific coping mechanisms, including problem solving, seeking social support, and avoidance. There was a correlation for greater reliance on avoidance with a greater increase in [reaction time](#) after night call.

"Reaction time increased considerably in anesthesiologists after a night call duty," the authors write. "Greater subjective reliance on avoidance as a coping strategy was associated with greater deterioration in performance."

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Anesthesiologist reaction time increased after night call (2016, December 12) retrieved 23 April 2024 from <https://medicalxpress.com/news/2016-12-anesthesiologist-reaction-night.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.