

Biomarker for oxidative stress plays a major role in hepatic inflammation

December 21 2016



A graphic illustration of the study. Credit: CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences

Diet-related diseases like non-alcoholic fatty liver disease (NAFLD) are known to have a major inflammatory component. However, the molecular pathways linking diet-induced changes with inflammation



remained elusive. In a new study, scientists at the CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences and the Medical University of Vienna identified crucial inflammatory processes in NAFLD. Moreover, the study published in *Hepatology* shows that malondialdehyde (MDA), a biomarker for oxidative stress, plays a key role in the development of NAFLD and can be neutralized by specific natural antibodies - a novel approach towards a potential therapy for the prevalent disease.

The combination of a nutrient rich, western diet and a lack of exercise is a lifestyle which can lead to serious health issues: Worldwide, the incidences of obesity, hypertension or insulin resistance are alarmingly high. As a consequence, risk of developing inflammation-associated diseases like type 2 diabetes, NAFLD and cardiovascular disease increased accordingly. However, the exact pathways that link the eating habits with the ensuing inflammation were so far not well understood.

The team of Christoph Binder, Professor of Atherosclerosis Research at the Medical University of Vienna and Principal Investigator at CeMM, in collaboration with Ronit Shiri-Sverdlov at the Maastricht University, Christoph Reinhardt at the University Medical Center of the Johannes Gutenberg University Mainz and the German Center for Cardiovascular Research Mainz was not only able to shed light on the biological processes that lead to the development of chronic inflammation upon a nutrient rich diet in mice. Moreover, the scientists found MDA to be a key player in hepatic inflammation which can be neutralized with natural antibodies. Their results were published in *Hepatology*.

The highly reactive molecule malondialdehyde, a product from lipid decomposition and biomarker for <u>oxidative stress</u>, accumulates on the surface of dying cells in the liver. Chemically bound to membrane proteins or phospholipids, they form so called MDA epitopes - Binder's research group showed that those MDA epitopes induce cytokine



secretion as well as leukocyte recruitment and thereby propagate existing inflammation and render it chronic.

"We were able to show in cell culture as well as in model organisms that those MDA epitopes play a major role in diet-induced hepatic inflammation," says Clara Jana-Lui Busch, one of the co-first authors of the study and PhD student at CeMM and the Medical University of Vienna. This was not the only insight of the study: "With intravenous injection of a specific MDA antibody which binds MDA epitopes selectively, we could ameliorate the inflammation in mice" Tim Hendrikx, the other co-first author and PostDoc in the group of Christoph Binder adds.

This study shows how the close collaboration of CeMM and the Medical University of Vienna fosters the development of a future precision medicine, says senior author Christoph Binder. "With cutting edge RNA sequencing methods and bioinformatic analyses of transcriptome data, we discovered key mechanisms in some of the most prevalent diseases and we confirmed those findings in mice models." Binder explains, and adds: "Above that, the administration of specific antibodies for MDA epitopes provide a promising new approach for the development of therapeutic strategies."

More information: Clara Jana-Lui Busch et al, Malondialdehyde epitopes are sterile mediators of hepatic inflammation in hypercholesterolemic mice, *Hepatology* (2016). <u>DOI:</u> <u>10.1002/hep.28970</u>

Provided by CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences



Citation: Biomarker for oxidative stress plays a major role in hepatic inflammation (2016, December 21) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2016-12-biomarker-oxidative-stress-major-role.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.