

New research suggests doctors' burnout should be treated as organisation-wide problem

December 6 2016

Current approaches to dealing with burnouts in doctors on an individual case-by-case basis is not effective and the issue should instead be tackled with organisation-wide initiatives, according to researchers at The University of Manchester and the University of Southampton.

A meta-analysis study, which brought together the results of previously conducted research, was carried out to explore the effectiveness of interventions in reducing [burnout](#) in [doctors](#). It explored the comparison between doctor-directed interventions that target the individual and organisation-directed interventions that target the working environment. The strength of the doctor's experience and the particular healthcare setting they worked in was also assessed.

The research concluded that while doctor-focused tactics such as mindfulness and cognitive behavioural are important, the greatest success at preventing and reducing burnout in doctors can be achieved through the adoption of organisation-directed approaches such as improved [working environment](#) and organisational culture.

Burnout is a major problem in the healthcare industry and is often driven by excessive workload, imbalance between job demands and skills, a lack of job control and prolonged stress. It is a syndrome consisting of emotional exhaustion, depersonalisation, and a diminished sense of personal accomplishment. Importantly, burnout can result in an increase

in medical errors, reduced quality of patient care, and lower patient satisfaction.

It was found that organisations that combined several elements such as structural changes, fostering communications between members of the health care team, and cultivating a sense of teamwork and job control tended to be the most effective in reducing burnout. However, such intense organisation-directed interventions were rare and had not been evaluated sufficiently.

What's more, the evidence indicated that young doctors starting out in their career, are at higher risk of burnout compared to those with more experience, and interventions focused on enhancing teamwork, mentoring, and leadership skills might be particularly suitable for this group.

Dr Maria Panagioti, Research fellow in Primary Care at the University of Manchester who led this study said: 'Our findings clearly show that we need more effective intervention models to prevent burnout in doctors. Such models could be organization-directed interventions which promote healthy individual-organization relationships and view burnout a problem of the whole healthcare systems.'

George Lewith, Professor of Health Research at the University of Southampton who supervised the research, said: "This work suggests that if we want to retain safe and professionally competent NHS clinicians working in very demanding front line jobs we need to support their mental and physical health and creating appropriate and enabling working environments for them. Efforts need to be focused on finding appropriate ways of reaching doctors who work in stressful environments to ensure their wellbeing is taken care of. If we don't patient safety could be at risk."

More information: Maria Panagioti et al. Controlled Interventions to Reduce Burnout in Physicians, *JAMA Internal Medicine* (2016). [DOI: 10.1001/jamainternmed.2016.7674](https://doi.org/10.1001/jamainternmed.2016.7674)

Provided by University of Manchester

Citation: New research suggests doctors' burnout should be treated as organisation-wide problem (2016, December 6) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-12-doctors-burnout-organisation-wide-problem.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.