

Exploring future preventive strategies for patients at imminent risk of hip fracture

December 19 2016

Hip fractures are of great concern as they are the most severe type of fracture in osteoporotic patients, associated with premature death, and commonly leading to long-term physical disability, impaired capacity to perform daily activities and live independently, and reduced quality of life. In the Western world, the lifetime risk for hip fracture in women over the age of 50 years is 18-23% and by 2050, the worldwide annual number of hip fractures is expected to reach 4.5 to 6.3 million, reflecting the continuous ageing of the population.

Prevention of [hip fracture](#) is currently based on pharmacological intervention together with [regular physical activity](#), prevention of falls, and correction of nutritional deficiencies, including vitamin D repletion. However, the efficacy of the various measures to reduce hip fractures is at most 50% and may need months or years before becoming effective. Additional challenges include the high treatment gap (even among those who have already suffered a hip fracture) and that 50% or more of patients who are on pharmacological treatment discontinue it within a year. Clearly there are unmet needs for patients at imminent risk of hip fractures that need to be addressed, including exploring potential new approaches.

A new review authored by a European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO)/International Osteoporosis Foundation (IOF) working group has looked at the presently available approaches for prevention of hip fracture and at potential future approaches to address unmet needs in

very high-risk patients.

The panel discussions concluded:

- There is a need to explore complementary approaches to immediately prevent hip fractures in very high risk patients, particularly following a first hip fracture.
- The challenges of current approaches include poor adherence and the lag time in onset of fracture risk reduction with medications.
- A possible role of orthopaedic interventions in strengthening bone should be explored and discussed.
- Procedures to test may include prophylactic nailing, femoroplasty with cement, and bone grafting with osteoconductive or osteoinductive materials.
- Clinical studies aimed at demonstrating the benefits and safety profile of surgical approaches are however challenging - including the need to use clear surrogate markers for bone strength improvement and identification of the level of high risk most likely to benefit from the procedure.
- The frail elderly patient, who is the most likely to be in need of, and benefit from, [fracture prevention](#) is also the most susceptible to comorbidities, and pre- and post-surgery complications.
- Target patient and intervention thresholds for surgical options remain to be established in the context of risk-benefit analysis - overall, the benefits should be in line with the risks.

Professor René Rizzoli, lead author and Chair of the ESCEO Scientific Advisory Board said: "Given the very serious, indeed life-threatening, repercussions of hip fractures, we must do all we can to find effective strategies for prevention, including in the frail elderly. Notwithstanding the many challenges involved, the presently untapped potential of preventive surgical interventions for patients at imminent risk of hip fracture deserves further investigation by researchers in the field."

More information: Serge Ferrari et al, Unmet needs and current and future approaches for osteoporotic patients at high risk of hip fracture, *Archives of Osteoporosis* (2016). [DOI: 10.1007/s11657-016-0292-1](https://doi.org/10.1007/s11657-016-0292-1)

Provided by International Osteoporosis Foundation

Citation: Exploring future preventive strategies for patients at imminent risk of hip fracture (2016, December 19) retrieved 13 March 2024 from <https://medicalxpress.com/news/2016-12-exploring-future-strategies-patients-imminent.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--