

# History of TBI linked to poor outcomes for those who are homeless, have mental illness

December 20 2016

---

Among homeless adults with mental illness, having a history of head injuries is associated with a greater risk of adverse health conditions, new research indicates.

Homeless adults with [mental illness](#) and a [history](#) of [traumatic brain injury](#) are also more likely to use emergency departments, report unmet health care needs and have contact with the criminal justice system than those without a history, according to a study published today in the *Journal of Psychiatric Research*.

This population also experienced an increased likelihood of being suicidal or having previously attempted suicide, the study said.

"Though it didn't prove a causal relationship, our research demonstrates that there is an association between specific mental health diagnoses and traumatic [brain injury](#)," said Dr. Stephen Hwang, lead author of the study and director of the Centre for Urban Health Solutions of St. Michael's Hospital.

"Head injuries are clearly a significant burden to people who are homeless or vulnerably housed, and we need to consider offering specialized services such as clinical treatment plans and supportive housing to help alleviate that burden."

Approximately 53 per cent of [homeless adults](#) with a mental illness had a reported history of traumatic brain injury, which is comparable to the

rate of reported traumatic brain injury in the general homeless population. Aboriginal participants were two times more likely than Caucasian participants to have a reported history of TBI.

All 2,088 study participants had a severe mental illness diagnosis with or without a diagnosis of a substance use disorder. Nearly all mental health and both alcohol and substance misuse diagnoses were associated with increased likelihood of a history of TBI in the study sample, with the exception of psychosis.

Although all participants had a mental illness, participants with specific mental illnesses were substantially more likely to have reported a history of TBI, most notably post-traumatic stress disorder (three times more likely than those without the disorder), panic disorder (2.5 times), depression (two times) and alcohol misuse disorder (two times).

"Earlier studies have looked at traumatic brain injury among people who experience homelessness who may or may not have had a mental health diagnosis. In comparison, our study included only individuals who were experiencing both homelessness and mental illness," said Agnes Gozdzik, a research associate in the Centre for Urban Health Solutions and a senior author of the study.

"By narrowing the diversity in the sample, we've shown that even among those with significant [mental health](#) issues, having a history of [head injuries](#) increases the risk of a number of negative outcomes."

This study involved data from more than 2,000 people enrolled across five cities in the Mental Health Commission of Canada's At Home/Chez Soi study, a research project that evaluated a "Housing First" approach, where people are first provided with a place to live, without preconditions such as sobriety or seeing a psychiatrist, and then provided additional supports and services as needed. The underlying principle is

that people are better able to move forward with their lives if they are first housed.

Provided by St. Michael's Hospital

Citation: History of TBI linked to poor outcomes for those who are homeless, have mental illness (2016, December 20) retrieved 9 April 2024 from

<https://medicalxpress.com/news/2016-12-history-tbi-linked-poor-outcomes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--