

Hormone-disrupting compound could provide new approach to malaria control

December 15 2016



A female *Anopheles gambiae* mosquito is feeding on blood. Credit: Emily Lund, Harvard

A chemical that disrupts biological processes in female mosquitoes may be just as effective as insecticides in reducing the spread of malaria,

according to a new study from Harvard T.H. Chan School of Public Health.

The study will be published online December 15, 2016 in *PLOS Pathogens*.

"As insecticide resistance is spreading, new intervention methods to control mosquitoes are urgently needed," said Flaminia Catteruccia, associate professor of immunology and infectious diseases at Harvard Chan School and co-senior author of the study. "Our study provides a new strategy based on the use of a non-toxic compound that prevents transmission of [malaria](#) parasites without killing the mosquito."

Malaria kills about 500,000 people every year. People can become infected when bitten by mosquitoes carrying the [malaria parasite](#), *Plasmodium falciparum*. Typical strategies to prevent transmission involve the use of insecticide-treated [bed nets](#) and insecticide spraying, but *Anopheles* mosquitoes—the type that transmits *P. falciparum* parasites—are rapidly becoming resistant to insecticides.

In the new study, researchers treated *Anopheles* [female mosquitoes](#) with a synthetic chemical called DBH (dibenzoylhydrazine) to see how it would impact their [biological processes](#). DBH mimics the action of the steroid hormone "20E" (20-hydroxyecdysone), which plays a key role in the reproductive cycle of the female mosquito.

Various aspects of the mosquitoes' life cycle were disrupted after treatment with DBH, the researchers found. The females produced and laid fewer eggs; they didn't mate successfully; and they died more rapidly than non-treated mosquitoes. The effects were greater the higher the DBH dose. DBH-treated [mosquitoes](#) were also less likely to be infected by malaria parasites.

Using DBH in either bed nets or in sprays was equally effective, the researchers found. They also noted that DBH compounds are not toxic to mammals, which would make them ideally suited for use in bed nets, where low toxicity is essential.

"The study demonstrates the importance of basic research on mosquito biology for developing new tools against malaria," said Caroline Buckee, co-senior author and assistant professor of epidemiology at Harvard Chan School.

More information: "Disrupting mosquito reproduction and parasite development for malaria control," Lauren M. Childs, Francisco Y. Cai, Evdoxia G. Kakani, Sara N. Mitchell, Doug Paton, Paolo Gabrieli, Caroline O. Buckee, Flaminia Catteruccia, *PLOS Pathogens*, online December 15, 2016, [DOI: 10.1371/journal.ppat.1006060](https://doi.org/10.1371/journal.ppat.1006060)

Provided by Harvard T.H. Chan School of Public Health

Citation: Hormone-disrupting compound could provide new approach to malaria control (2016, December 15) retrieved 19 April 2024 from <https://medicalxpress.com/news/2016-12-hormone-disrupting-compound-approach-malaria.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--