


Immune system cells cause severe malaria complication in mouse brain

December 1 2016



The brain image (ECM brainstem pathology) shows severe vascular leakage in the brainstem of mouse with fatal cerebral malaria. This type of pathology is induced by cerebral herniation (or, pressure placed on the brainstem due to brain swelling). The white in this image represents a vascular dye called Evan's blue. The blue color in this image represent cell nuclei. Credit: The image was generated in Dr. McGavern's laboratory at the National Institute of Neurological Disorders & Stroke by Dr. Phillip Swanson II.

Immune system cells known as cytotoxic T cells attack blood vessels and cause fatal swelling in the brains of mice with a condition that mirrors a


severe complication of malaria in humans. These are the findings of a new study published in *PLOS Pathogens*.

After contracting malaria, some people develop a complication known as cerebral malaria, which causes brain swelling and bleeding. Cerebral malaria kills 15 to 30 percent of the people it afflicts, but the mechanisms that cause the disease are not completely understood.

To help clarify the causes of cerebral malaria, Dr. Phillip Swanson of the National Institutes of Health, Maryland, and colleagues peered into the brains of mice infected with a parasite that causes a similar fatal condition. They used a technique called intravital microscopy to capture movies of immune cell activity in the living mouse brain as the disease progressed.

The research revealed that cytotoxic T [cells](#) aggregated along blood vessels in the diseased mouse brain, interacting with tiny parasite fragments presented by the vessels. These interactions were directly associated with vascular leakage. The leaking blood vessels caused the brain to swell, and the resulting pressure killed neurons that keep the heart and lung functioning.

When the scientists injected the mice with antibodies that prevented T cells from sticking to blood vessels, the fatal swelling was prevented. This confirmed that T cells play a major role in causing the mouse version of cerebral malaria. It also suggests potential new treatment strategies, should further research confirm that a similar mechanism is responsible for cerebral malaria in humans.



This image (intravascular localization of parasite-specific T cells) captured by two-photon microscopy through the skull bone of mouse with fatal cerebral malaria. It shows parasite-specific CD8 T cells (green) attacking the vascular lumen of cerebral blood vessels. These interactions are cause fatal brain swelling and cerebral herniation. Blood vessels in this image are shown in red. Credit: The image was generated in Dr. McGavern's laboratory at the National Institute of Neurological Disorders & Stroke by Dr. Phillip Swanson II.

"By watching immune cells function in the living brain during [cerebral malaria](#), the investigators of this study revealed that parasite-specific CD8 T cells attack the wall of the cerebral [blood vessels](#), causing excessive swelling and damage to a vital brain center. This fatal disorder can be prevented by therapeutically displacing the pathogenic T cells from the vessel wall."

More information: Swanson PA, II, Hart GT, Russo MV, Nayak D, Yazew T, Peña M, et al. (2016) CD8+ T Cells Induce Fatal Brainstem Pathology during Cerebral Malaria via Luminal Antigen-Specific Engagement of Brain Vasculature. *PLoS Pathog* 12(12): e1006022. [DOI: 10.1371/journal.ppat.1006022](#)

Provided by Public Library of Science

Citation: Immune system cells cause severe malaria complication in mouse brain (2016, December 1) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-12-immune-cells-severe-malaria-complication.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--