

## New test to identify obese women at high risk of developing diabetes in pregnancy

December 8 2016

In 2014, around seven million women in the UK were classified as obese. By 2025, it is expected to affect 1 in 5 women in the world. Obesity is a major risk factor for gestational diabetes, increasing the likelihood of the disorder three - five fold. Women with the disorder require intensive antenatal care to control blood glucose and to identify other common complications, particularly fetal macrosomia - a newborn who's significantly larger than average.

In practice today, all obese pregnant <u>women</u> are categorised as being of equally high risk of gestational diabetes, whereas in reality, only around 25% will develop the disorder.

In the study, funded by the National Institute for Health Research (NIHR), the team looked at how to correctly identify obese women with heightened risk, early in pregnancy, and as a result, enable timely targeted intervention to those women most likely to benefit.

From the many factors tested, those that predicted gestational diabetes included older age, disease in a previous pregnancy, higher blood pressure and anthropometric measures such as skin thicknesses, waist and mid-arm circumferences. A number of blood tests, such as Haemoglobin A1c also added strength to the predictive tool.

Out of the 1303 women in the study, 337 were affected by gestational diabetes.



Lead author, Dr Sara White from King's College London said: "There is currently no accepted strategy to identify <u>obese women</u> at high risk of gestational diabetes, early in pregnancy. Today, all those classified as obese are considered <u>high risk</u>. With escalating rates of obesity worldwide, a more accurate way of defining risk is necessary in this group.

"In this, the largest and most comprehensive study to date, we have used an extensive range of different measures to develop prediction tools. One of our models focused on a few clinical factors and biomarkers already readily available in clinical practice and which incurred minimal cost. In addition, we have identified a model that does not require blood sampling, which could be developed for low and <u>middle income</u> <u>countries</u> where the prevalence of gestational diabetes and obesity is rapidly increasing.

"Clinical use of these tests would enable prompt intervention and correctly target those at highest risk and therefore most likely to benefit."

Dr White is continuing her research into <u>gestational diabetes</u> with the support of Diabetes UK.

Provided by King's College London

Citation: New test to identify obese women at high risk of developing diabetes in pregnancy (2016, December 8) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2016-12-obese-women-high-diabetes-pregnancy.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.