

## Oxidative stress tied to early-onset androgenetic alopecia

December 27 2016



(HealthDay)—Younger patients with early-onset androgenetic alopecia



(AGA) have increased oxidative stress, according to a study published online Dec. 16 in the *Journal of Cosmetic Dermatology*.

Hilal Kaya Erdogan, M.D., from Eskischir Osmangazi University in Turkey, and colleagues evaluated the oxidative stress status in 33 male patients with early-onset AGA and 30 men without AGA between ages of 18 and 30 years old. Measurements included total oxidant levels (TOS), total antioxidant levels (TAS), and oxidative <u>stress</u> index (OSI).

The researchers observed no difference for TAS level between the patient and control groups. However, TOS and OSI were significantly higher in the patient group. There was a highly significant negative correlation between TAS level and both age and disease duration. However, when TAS, TOS, and OSI levels were evaluated by AGA stage, there was no significant difference between groups. OSI level was significantly higher in patients with a family history of AGA.

"There is need for further molecular studies on the role of <u>oxidative</u> <u>stress</u> in the etiopathogenesis of AGA," the authors write.

**More information:** <u>Full Text (subscription or payment may be</u> <u>required)</u>

Copyright © 2016 HealthDay. All rights reserved.

Citation: Oxidative stress tied to early-onset androgenetic alopecia (2016, December 27) retrieved 17 May 2024 from <u>https://medicalxpress.com/news/2016-12-oxidative-stress-tied-early-onset-androgenetic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.