

Psychiatric conditions linked to increased risk of long-term opioid use

December 29 2016

A wide range of pre-existing psychiatric and behavioral conditions and the use of psychoactive drugs could be important risk factors leading to long-term use of opioid pain medications, reports a study in *Pain*, the official publication of the International Association for the Study of Pain (IASP).

Using a nationwide insurance database, the researchers identified 10.3 million [patients](#) who filed insurance claims for opioid prescriptions between 2004 and 2013. The study looked at whether pre-existing psychiatric and behavioral conditions and use of psychoactive medications were predictors of later opioid use.

"We found that pre-existing psychiatric and behavioral conditions and psychoactive medications were associated with subsequent claims for [prescription opioids](#)," write Patrick D. Quinn, PhD, of Indiana University, Bloomington, and colleagues. The association appears stronger for long-term opioid use, and especially for patients with a previous history of substance use disorders.

The results also suggest that some outcomes viewed as harmful outcomes of opioid use—substance use disorders, depression, suicidal or self-injuring behavior, and motor vehicle crashes—are also *predictors* of which patients are at risk of long-term use of prescription opioids.

Increased Rates of Opioid Prescribing for People with

Mental Health Conditions

Overall, the results suggested a "modest" increase in any opioid prescriptions for patients with previous psychiatric or behavioral conditions (depression or anxiety disorders, opioid or other substance use disorders, suicide attempts or other self-injury, motor vehicle crashes, and sleep disorders) or use of psychoactive medications.

About 1.7 percent of patients with opioid prescriptions become long-term opioid users (six months or longer). But the risk became substantially higher for patients with [mental health conditions](#) or psychoactive medication use. Relative increases in rates of long-term opioid use ranged from 1.5 times for patients taking medications for attention-deficit/hyperactivity disorder, to about 3 times for those with previous [substance use disorders](#) other than opioids, to nearly 9 times for those with previous opioid use disorders.

Amid the continuing opioid epidemic, it's important to understand which patients select (or are selected for) treatment with these pain medications. Previous studies have suggested a pattern of "adverse selection": patients at greatest risk of harmful outcomes, including those with substance abuse and other psychiatric conditions, may be more likely to be prescribed opioids in higher doses and for longer durations.

"Our results add to existing evidence that the risk of long-term opioid receipt associated with [pre-existing] psychiatric and behavioral conditions is widespread and relates to multiple diagnoses and [psychoactive medications](#)," Dr. Quinn and coauthors write. The results add to previous evidence suggesting that opioids are more likely to be prescribed to certain groups of patients at high risk for adverse outcomes.

Dr. Quinn and coauthors conclude: "Our findings support the ideas that

clinical practice has deviated from the 'careful selection' under which most clinical trials are conducted and that thorough [mental health](#) assessment and intervention should be considered in conjunction with the use of long-term opioid therapy."

More information: Patrick D. Quinn et al. Incident and long-term opioid therapy among patients with psychiatric conditions and medications, *PAIN* (2017). [DOI: 10.1097/j.pain.0000000000000730](https://doi.org/10.1097/j.pain.0000000000000730)

Provided by Wolters Kluwer Health

Citation: Psychiatric conditions linked to increased risk of long-term opioid use (2016, December 29) retrieved 9 April 2024 from <https://medicalxpress.com/news/2016-12-psychiatric-conditions-linked-long-term-opioid.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--