


Single-dose LMWH can interfere with algorithm for DVT diagnosis

December 22 2016



(HealthDay)—A single dose of low-molecular-weight heparin (LMWH)

given the previous day can reduce D-dimer and potentially risk non-identification of a deep vein thrombosis (DVT), according to a research letter published online Dec. 15 in the *International Journal of Laboratory Hematology*.

Peter M. Baker and David M. Keeling, M.B., Ch.B., from the Oxford Haemophilia and Thrombosis Centre in the United Kingdom, examined whether a single dose of LMWH given in the community when a patient presented to their general practitioner too late in the day for referral would render the D-dimer algorithm unusable the following morning. Paired samples were collected for D-dimer estimation from 50 patients pre-administration of LMWH and on presentation to the DVT clinic the following morning.

The researchers found that the sample taken post-administration of a single dose of LMWH showed a significantly lower D-dimer, from 2,670 $\mu\text{g/L}$ to 2,274 $\mu\text{g/L}$ (16.4 percent reduction; $P < 0.01$). Reduction in D-dimer crossed the normal range of

Citation: Single-dose LMWH can interfere with algorithm for DVT diagnosis (2016, December 22) retrieved 3 May 2024 from <https://medicalxpress.com/news/2016-12-single-dose-lmwh-algorithm-dvt-diagnosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.