

New findings about stem cells in the brain of patients with epilepsy

December 15 2016



Generalized 3 Hz spike and wave discharges in a child with childhood absence epilepsy. Credit: Wikipedia.

Neural stem cells have been found in epileptic brain tissue—outside the regions of the brain where they normally reside. In a group of patients who underwent surgery for epilepsy, over half had stem cells where healthy individuals do not have them, according to a study from Sahlgrenska Academy.

"We have confirmed what earlier studies indicated, and gained new knowledge about [molecular characteristics](#) of these [neural stem cells](#)," says Milos Pekny, professor at the Institute of Neuroscience and Physiology.

Neural stem cells with the ability to form new neurons in the brain are normally present in the hippocampus (the part of the brain connected to learning and memory) and in the subventricular zone of the brain. However, in 8 of 14 patients in the study, neural stem cells were present outside these regions, namely in the epileptic [brain tissue](#) that was surgically removed.

Over 50 operations of this kind are conducted in Sweden each year, of which about 20 in Gothenburg, with most patients becoming seizure-free or having significant reduction in seizure frequency. After an extensive investigation aiming at the precise localization of the epileptic focus, brain tissue that is damaged due to malformation, injury or other cause, is surgically removed.

Good teamwork

Surgeons, neurologists and neuroscientists worked side by side in the current study, which followed ethical approval and informed consent from each patient. For research purposes, the team was allowed to examine a small part of the removed tissue used for histopathological examination - in the operating room and in the research laboratory, just several minutes after removal.

"About 60% of the patients had epileptic tissue that contained neural stem cells that could be converted into neurons, astrocytes and oligodendrocytes (the three types of [brain cells](#) that neural stem cells can differentiate into) when they were later grown in the laboratory," says Milos Pekny.

"This may point to a greater plasticity in the epileptic tissue, which to some extent can be compared to the brain tissue of a newborn," continues Milos Pekny.

Scientists have gained a better molecular understanding of the region of the brain in individuals with epilepsy which - due to a developmental abnormality, trauma, stroke, or a growing tumor - has stopped responding to control signals, and this results in recurrent seizures.

Improved understanding

"The knowledge gained in this study primarily helps to improve our understanding of the brain responses in epilepsy," outlines Milos Pekny.

Scientists have long speculated that astrocytes, the cell type that controls many neuronal functions, give rise to neural [stem cells](#) in damaged brain tissue.

"Our study suggest that this is not the case, at least in epilepsy, and it contributes the advancement of our understanding of what can happen in the [brain](#) in people with epilepsy," says Milos Pekny.

More information: Neural Progenitor Cells in Cerebral Cortex of Epilepsy Patients do not Originate from Astrocytes Expressing GLAST.
[DOI: 10.1093/cercor/bhw338](https://doi.org/10.1093/cercor/bhw338)

Provided by University of Gothenburg

Citation: New findings about stem cells in the brain of patients with epilepsy (2016, December 15) retrieved 25 April 2024 from <https://medicalxpress.com/news/2016-12-stem-cells-brain-patients-epilepsy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.