

## Tracking down therapy-resistant leukaemia cells

December 15 2016



Leukemia cells finding shelter against the cancer therapy. Credit: HMGU



Dr. Irmela Jeremias from Helmholtz Zentrum München and her colleagues have succeeded in finding a small population of inactive leukaemia cells that is responsible for relapse of the disease. Now the way is paved for research into new therapies that prevent disease relapse by eliminating the remaining, so-called dormant leukaemia cells. The research results have now been published in the *Cancer Cell* journal.

Chemotherapy often fails in leukaemia when <u>resistant cells</u> survive treatment and bring about a <u>relapse</u> of the disease (recurrence). New therapies are therefore needed to eliminate these cells. The team of scientists headed by Jeremias, head of the 'Apoptosis' research group in the Gene Vectors Research Unit (AGV) at the Helmholtz Zentrum München, has now isolated and characterized therapy-resistant cells for the first time. "Previously the biological principles responsible for a relapse in leukaemia were not fully understood," says Jeremias. "Our new approach is to isolate dormant cells, which gives us the first possibility of developing therapies that switch off these cells."

## Isolated cells respond to drugs

"We have found a method to dissociate dormant leukaemia cells from their surroundings, where they are safe from attacks by therapeutics," explains Sarah Ebinger, <u>doctoral candidate</u> in the AGV and the article's first author. With the help of modern genetic engineering and dyes that mark cell growth, the scientists isolated cells and identified a rare cell type that resembled cells triggering relapse. These cells were inactive and resistant to therapy. "We then found out that these cells, once they have been dissolved out of their surroundings, are indeed susceptible to therapy and react well to therapeutics," adds Erbey Özdemir, doctoral candidate in the AGV. "This has brought us a small step closer to the global goal of preventing disease relapse in patients suffering leukaemia," says Jeremias. "It might serve as basis for new therapies that destroy resistant leukaemia <u>cells</u> before they induce relapse."



**More information:** Sarah Ebinger et al. Characterization of Rare, Dormant, and Therapy-Resistant Cells in Acute Lymphoblastic Leukemia, *Cancer Cell* (2016). <u>DOI: 10.1016/j.ccell.2016.11.002</u>

## Provided by Helmholtz Association of German Research Centres

Citation: Tracking down therapy-resistant leukaemia cells (2016, December 15) retrieved 5 June 2024 from <u>https://medicalxpress.com/news/2016-12-tracking-therapy-resistant-leukaemia-cells.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.