


Risk of uterine fibroids found to be lower in women using statins

December 21 2016



(HealthDay)—The use of statins is associated with a lower risk of

uterine fibroids and fibroid-related symptoms, according to a study published in the December issue of the *American Journal of Obstetrics & Gynecology*.

Mostafa A. Borahay, M.D., Ph.D., from the University of Texas Medical Branch at Galveston, and colleagues examined the association between statin use and the risk of [uterine fibroids](#) and fibroid-related symptoms in commercially insured women diagnosed with hyperlipidemia. The authors matched 47,713 women with diagnosed uterine fibroids with 143,139 controls without fibroids (1-to-3 ratio).

The researchers found that exposure to statins within two years prior to the event/index date was associated with a decreased risk of uterine fibroids (odds ratio [OR], 0.85). In subanalysis of the women with uterine fibroids, statin users had a lower likelihood of having menorrhagia (OR, 0.88), anemia (OR, 0.84), or pelvic pain (OR, 0.85) and of undergoing myomectomy (OR, 0.76) compared with nonusers of [statins](#).

"Further studies, including [randomized controlled trials](#), may be warranted," the authors write.

One author disclosed financial ties to AbbVie.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Risk of uterine fibroids found to be lower in women using statins (2016, December 21) retrieved 23 April 2024 from <https://medicalxpress.com/news/2016-12-uterine-fibroids-women-statins.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.