

Breast cancer prognosis of African-American patients may improve with chemotherapy before surgery

January 19 2017, by Latina Emerson



Credit: Georgia State University

Administering chemotherapy to African-American breast cancer patients prior to surgery could improve their prognosis and survival rates from

the disease, according to a new study.

A research team led by Georgia State University found African-American breast cancer patients who receive chemotherapy prior to surgery exhibit trends of reduced regional (in lymph nodes) recurrence and distant (in remote organs, such as the liver, lungs and brain) recurrence of tumors, which may help diminish the inequality in breast cancer-related clinical outcomes between African-American and European-American patients.

There is a significant disparity in [breast cancer mortality](#) between African-American and European-American breast cancer patients. Despite a similar number of reported cases of breast cancer among African-American and European-American women, African-Americans experience a more aggressive clinical course and a 40 percent higher death rate than European-Americans among premenopausal and menopausal breast cancer patients.

Recurrent breast cancer has hindered the successful management of this disease for decades and is one of the primary factors for the racial disparity in prognosis and outcomes. Differences in recurrence rates and patterns between the races following various forms of treatment have not been thoroughly investigated.

This is the first clinical study to suggest that neoadjuvant chemotherapy (treatment prior to surgery) may improve [breast cancer recurrence](#) rates and patterns in African-Americans. The results are published in the journal *PLOS ONE*.

Researchers at Georgia State analyzed clinical data from a large cohort of breast cancer patients treated at Northside Hospital in Atlanta from 2005 to 2015. The patients' self-reported races were primarily African-American and European-American. The researchers studied rates and

patterns of [tumor recurrence](#) after hormone, radiation and chemotherapy among African-American and European-American breast cancer patients.

"We found that, in general, African-American breast cancer patients exhibit increased likelihood for tumor recurrence, particularly to regional and distant sites, after receiving any combination of adjuvant therapy (treatment following surgery) compared to European-American breast [cancer patients](#). This higher incidence of tumor recurrence can contribute to a poorer prognosis," said Nikita Wright, first author of the study and a senior Ph.D. student in Dr. Ritu Aneja's laboratory in Georgia State's Biology Department.

When [breast cancer](#) recurs, it is more challenging to treat regional and distant tumors than local tumors (in the breast), Wright explained.

"Interestingly, we found that neoadjuvant chemotherapy actually reversed these recurrence trends," Wright said. "We found that African-American [breast cancer patients](#) responded better to neoadjuvant chemotherapy than European-American patients. Among patients who received [neoadjuvant chemotherapy](#), African-Americans exhibited trends of lower regional and distant tumor recurrence than European-Americans, but higher local recurrence, which is easier to manage clinically and is associated with a relatively better prognosis."

More information: Nikita Wright et al. Distinctions in Breast Tumor Recurrence Patterns Post-Therapy among Racially Distinct Populations, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0170095](https://doi.org/10.1371/journal.pone.0170095)

Provided by Georgia State University

Citation: Breast cancer prognosis of African-American patients may improve with chemotherapy before surgery (2017, January 19) retrieved 20 April 2024 from <https://medicalxpress.com/news/2017-01-breast-cancer-prognosis-african-american-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.