

## Timing of chemo affects inflammation, mice study suggests

January 24 2017, by Jeff Grabmeier



Breast cancer cells. Image: Annie Cavanagh

The time of day that breast cancer chemotherapy drugs are given affects the amount of damaging inflammation in the body, a new study in mice suggests.

That's important because inflammation, particularly in the brain, is believed to contribute to many of the neurological <u>side effects</u> of chemotherapy in humans, such as depression, anxiety and short-term memory loss.

"Timing of <u>drug</u> administration can have a big effect on inflammation, and that may potentially affect a wide variety of harmful side effects," said Courtney DeVries, co-author of the study and professor of neuroscience at The Ohio State University's Wexner Medical Center.



But the results also showed an important complicating factor: The inflammatory effects were opposite in the brain versus the spleen depending on the time the drugs were given.

"The spleen and the brain were out of phase. When inflammation was high in one, it was low in the other," said co-author Randy Nelson, also a professor of neuroscience at Ohio State's Wexner Medical Center.

The study was published today in the journal Scientific Reports.

These results mark another step in how scientists understand cancer chronotherapy—using the body's daily circadian rhythms to time treatments so they are most effective and have the fewest side effects.

"I think we know enough about circadian rhythms in terms of physiology that we can start translating these findings into medical research and practice in humans," Nelson said.

The research team studied the drugs cyclophosphamide and doxorubicin, a common treatment for breast cancer, in female <u>mice</u>. The mice did not have tumors because this study was designed just to look at the inflammatory response to the drugs.

The mice were injected with the drugs two hours after daylight (which is their inactive period) or two hours after lights were turned off (their active period).

Researchers collected tissue and checked for signs of inflammation in the spleen, an important immune system organ. The also looked at two sites in the brain: the hypothalamus and the hippocampus. Tissue was collected at 1, 3, 9 or 24 hours after the injection.

Results showed that injecting mice with the chemotherapy drugs in their



inactive phase (daylight) increased the expression of genes that promoted inflammation within the spleen. The researchers found increased production of two toxic drug metabolites, products of the chemo drugs. These metabolites are related to inflammation and one of them, doxorubicinol, causes heart damage in some patients.

"The timing of when patients receive drugs has the potential to reduce serious side effects," DeVries said.

The effect was different for mice that received the drugs two hours after the lights were out. In these mice, inflammation measured in the spleen was much lower than the mice injected during the day.

The pattern was reversed in the brain, however. Mice injected at night had increases in pro-inflammatory gene expression, but showed less evidence of inflammation when injected during the day.

The difference in <u>inflammation response</u> between the brain and the rest of the body needs to be better appreciated by physicians, Nelson said.

"Many times physicians measure markers of <u>inflammation</u> in the blood and assume that is what is happening in the brain, but we found that just the opposite is happening," he said.

Why is there a difference between how the brain and spleen are reacting to these chemotherapy drugs?

Other research has shown that the clock genes in the <u>brain</u> are out of sync with those in other organs. Expression of these clock genes may affect how the drugs are processed, or metabolized, in the body, Nelson said.

These findings can't tell us the best time to administer <u>chemotherapy</u>



drugs in humans, DeVries said of the new study.

"But it does demonstrate that time of day is an important factor that needs to be considered in when the drugs are given," she said.

Nelson noted that this study only looked at administering drugs at two time points.

"There may be a sweet spot that maximizes the efficacy of the drugs and minimizes the side effects. We don't know that yet."

Provided by The Ohio State University

Citation: Timing of chemo affects inflammation, mice study suggests (2017, January 24) retrieved 21 May 2024 from <u>https://medicalxpress.com/news/2017-01-chemo-affects-inflammation-mice.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.