

Experts urge for wider prescription of statins in treatment and prevention

January 18 2017

World-renowned researchers from the Charles E. Schmidt College of Medicine at Florida Atlantic University as well as Harvard Medical School address the possible but unproven link between statins and diabetes, as well as the implications of prescription of statins for clinicians and their patients, in a commentary published in the prestigious *American Journal of Medicine*. The editor-in-chief of the journal published the commentary and an editorial he wrote online ahead of print.

Charles H. Hennekens, M.D, Dr.P.H., the first Sir Richard Doll professor and senior academic advisor to the dean, the Charles E. Schmidt College of Medicine at FAU; Bettina Teng, BA, a recent pre-med honors graduate of the Harriet L. Wilkes Honors College at FAU; and Marc A. Pfeffer, M.D., Ph.D., the Dzaou professor of medicine at HMS, emphasize to clinicians that the risk of diabetes, even if real, pales in comparison to the benefits of [statins](#) in both the treatment and [primary prevention](#) of heart attacks and strokes.

"The totality of evidence clearly indicates that the more widespread and appropriate utilization of statins, as adjuncts, not alternatives to therapeutic lifestyle changes, will yield net benefits in the treatment and primary prevention of heart attacks and strokes, including among high, medium and low risk patients unwilling or unable to adopt [therapeutic lifestyle changes](#)," said Hennekens.

In the accompanying editorial, Joseph S. Alpert, M.D., editor-in-chief

and a renowned cardiologist and professor of medicine at the University of Arizona School of Medicine, reinforces these important and timely clinical and public health challenges in treatment and primary prevention.

"There is no threshold for low density lipoprotein cholesterol below which there are no net benefits of statins either in the treatment or primary prevention of heart attacks and strokes," said Alpert.

The authors and editorialist express grave concerns that there will be many needless [premature deaths](#) as well as preventable heart attacks and strokes if patients who would clearly benefit from statins are not prescribed the drug, refuse to take the drug, or stop using the drug because of ill-advised adverse publicity about benefits and risks, which may include misplaced concerns about the possible but unproven small risk of diabetes.

"These public health issues are especially alarming in primary prevention, particularly among women, for whom cardiovascular disease also is the leading cause of death, and for whom there is even more underutilization of statins than for men," said Hennekens.

At its national meeting in November 2013, the American Heart Association, in collaboration with the American College of Cardiology, presented and published its new guidelines for the use of statins in the treatment and primary prevention of heart attacks and strokes, in which the organizations also recommended wider utilization in both treatment and prevention.

According to the United States Centers for Disease Control and Prevention, heart disease is the leading killer among men and women, causing approximately 600,000 deaths each year.

Provided by Florida Atlantic University

Citation: Experts urge for wider prescription of statins in treatment and prevention (2017, January 18) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-01-experts-urge-wider-prescription-statins.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.