

Glucose supplementation significantly reduces length of induced labor in childbirth

January 23 2017

In a study to be presented Thursday, Jan. 26, in the oral concurrent session, at the Society for Maternal-Fetal Medicine's annual meeting, The Pregnancy Meeting, researchers with the Université de Sherbrooke in Sherbrooke, Quebec, Canada will present their findings in a study titled, Reduction of total labor length through the addition of parenteral dextrose solution in induction of labor in nulliparous: results of DEXTRONS prospective randomized controlled trial. The study investigated the use of glucose to shorten induced labor in nulliparous (first time giving birth). The primary outcome studied was the total length of active labor.

Prolonged [labor](#) can be harmful to maternal and fetal health. Few medical interventions are known to shorten labor duration. Because muscle performance is known to be improved by [glucose](#) supplementation, the researchers tested whether adding glucose to the intravenous hydration solution women receive during labor could accelerate labor. Two hundred pregnant women were randomly assigned to receive either a standard hydration solution containing salt and water or a solution containing glucose, salt and water.

Josianne Pare, M.D., with the Department of Obstetrics & Gynecology at the University of Sherbrooke and the presenter of the research at the SMFM annual meeting, explained, "We found that the median duration of labor was 76 minutes shorter in the group of women receiving glucose. There was no difference in the mode of delivery (cesarean section, forceps, etc.), or the neonatal well-being measures." Pare

continued, "Glucose supplementation therefore significantly reduces the total length of labor without increasing the rate of complication. This is great news for women experiencing induced labor."

The researchers concluded that, given the low-cost and safety of this intervention, glucose should be the solute of choice during labor.

More information: Abstract #32 Reduction of total labor length through the addition of parenteral dextrose solution in induction of labor in nulliparous: Results of DEXTRONS prospective randomized controlled trial , The Pregnancy Meeting, 2017.

Provided by Society for Maternal-Fetal Medicine

Citation: Glucose supplementation significantly reduces length of induced labor in childbirth (2017, January 23) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-01-glucose-supplementation-significantly-length-labor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.