

Inflammatory markers prognostic in primary sinonasal cancer

January 10 2017



(HealthDay)—Pretreatment neutrophil-to-lymphocyte ratio (NLR) and

platelet-to-lymphocyte ratio (PLR) can predict survival for patients with primary sinonasal cancers (SNCs), according to a study published online Dec. 29 in *Head & Neck*.

Mario Turri-Zanoni, M.D., from the University of Insubria in Varese, Italy, and colleagues conducted a [retrospective review](#) of patients with SNC who had been treated using endoscopic approaches from 2002 to 2014. They examined the value of NLR and PLR in predicting [recurrence](#) and mortality for 215 patients.

The researchers found that in analysis of [epithelial tumors](#) (adenocarcinoma and carcinoma) and advanced-stage tumors (pT3-T4), patients with higher NLR and PLR had shorter overall and disease-free survival. NLR and PLR were independent prognostic factors for disease-free survival, with the risk of recurrence reduced among [patients](#) with NLR

Citation: Inflammatory markers prognostic in primary sinonasal cancer (2017, January 10) retrieved 5 May 2024 from <https://medicalxpress.com/news/2017-01-inflammatory-markers-prognostic-primary-sinonasal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
