

Pre-transplant depression may impair post-HCT outcomes

January 30 2017



(HealthDay)—For adult patients undergoing hematopoietic cell

transplantation (HCT), pre-transplant depression is associated with worse post-transplantation outcomes, according to a study published online Jan. 19 in *Cancer*.

Areej El-Jawahri, M.D., from the Massachusetts General Hospital in Boston, and colleagues analyzed data from the Center for International Blood and Marrow Transplant Research (CIBMTR) to compare outcomes for [adult patients](#) with hematologic malignancies with and without depression after autologous (3,786 [patients](#)) or allogeneic (7,433 patients) HCT.

The researchers found that among patients who underwent allogeneic HCT between 2008 and 2012, 15 and 85 percent, respectively, had pre-transplant depression and no depression. Pre-transplant depression correlated with lower overall survival (hazard ratio, 1.13; 95 percent confidence interval [CI], 1.04 to 1.23; P = 0.004) and with increased incidence of grade 2 to 4 acute graft-versus-host disease (hazard ratio, 1.25; 95 percent CI, 1.14 to 1.37; P hospital (means ratio, 0.98; 95 percent CI, 0.97 to 0.99; P = 0.002).

"Patients with pre-transplant [depression](#) represent a population that is at risk for post-transplant complications," the authors write.

The CIBMTR was partially funded by several pharmaceutical companies.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Pre-transplant depression may impair post-HCT outcomes (2017, January 30) retrieved

27 April 2024 from

<https://medicalxpress.com/news/2017-01-pre-transplant-depression-impair-post-hct-outcomes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.