

# Evaluation of recombinant antithrombin versus placebo in preterm preeclampsia

January 23 2017

---

In a study to be presented Friday, Jan. 27, in the late breaking oral session, at the Society for Maternal-Fetal Medicine's annual meeting, The Pregnancy Meeting, researchers with The PRESERVE-1 Study Group University of Texas Health Science Center at Houston—McGovern Medical School, Houston, Texas, and Yale School of Medicine, New Haven, Connecticut, present findings of a study titled Randomized double-blind placebo controlled evaluation of the safety and efficacy of recombinant Antithrombin versus placebo in preterm preeclampsia. The study was sponsored by rEVO Biologics, Inc.

Preeclampsia is a major cause of maternal and perinatal mortality and morbidity. In addition, women who develop early onset preeclampsia have markedly increased rate of complications both acute and long term such as cardiovascular disease, stroke, renal injury and metabolic syndrome. Infants born at less than 30 weeks have significant neonatal complications with prolonged stays in the [neonatal intensive care](#) unit. In addition, they are at increased risk for [chronic lung disease](#), cerebral palsy and other neurologic deficits. There are approximately 8,000 cases of early onset preeclampsia each year in the U.S. The estimated maternal and neonatal cost of these pregnancies is almost \$1.5 billion.

This trial investigated the effects of recombinant antithrombin (ATryn), a man-made version of antithrombin, a protein molecule found in blood that is produced by the liver, regulates the coagulation system, and has anti-inflammatory properties. It was studied to determine its potential to prolong gestation and improve maternal and neonatal outcomes. The

study's design was remarkable in that it was the largest randomized, controlled trial ever to be completed in patients who developed preeclampsia very early in pregnancy, 23-30 weeks' gestational age.

Baha Sibai, M.D. with the University of Texas Health Science Center at Houston and the presenter of the study at the SMFM annual meeting, reported "The results found no improvement in outcomes with such therapy. There were no reported safety events related to Recombinant Antithrombin" Future studies should investigate different novel targeted therapies to improve outcome in such pregnancies."

**More information:** Abstract LB02: Randomized double-blind placebo controlled evaluation of the safety and efficacy of recombinant Antithrombin versus placebo in preterm preeclampsia, The Pregnancy Meeting, 2017.

Provided by Society for Maternal-Fetal Medicine

Citation: Evaluation of recombinant antithrombin versus placebo in preterm preeclampsia (2017, January 23) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-01-recombinant-antithrombin-placebo-preterm-preeclampsia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--