


Synaesthesia 'hearing-motion' phenomenon more common than previously thought, says study

January 19 2017, by George Wigmore



Credit: AI-generated image ([disclaimer](#))

A little-known synaesthetic 'hearing-motion' phenomenon in which people hear faint sounds following totally silent visual flashes may be more common than previously realised, according to a new study from City, University of London.

The paper, which is published in the journal *Consciousness and Cognition*, is the first to show that a surprisingly high proportion of individuals report such sensations, and that these visually-evoked sounds seem to interfere with the ability to hear real sounds, with the flashes essentially having a deafening effect on individuals.

It is thought it may arise as humans are better at recording sound than images, therefore the ability to recode visual signals as sounds would take advantage of such abilities.

Most synaesthesia is rare, where stimulation of one sensory pathway leads to automatic, involuntary experience of a second sensory pathway, occurring in around 4 percent of the population. However, the new City study found that such 'hearing-motion' synaesthesia occurred in 22 percent of people when tested and it is thought that they may occur subliminally, disrupting detection of real auditory signals.

The frequent natural correlation between visual and auditory stimuli might explain the surprising prevalence of this phenomenon. Overall, the results suggest that learned correspondences between strongly correlated senses may provide a precursor for some synaesthetic abilities.

The research supports the idea that in fact some forms of synaesthesia might be grounded on normal mechanisms involved in forming and reinforcing associations between different senses, and that such synaesthetic associations might form more frequently when they occur between senses which are naturally related, such as sound and vision, as they can be consistently reinforced.

To investigate the effect, 40 randomly-sampled participants with normal hearing and vision were presented with pairs of either visual or auditory Morse-code like patterns, and had to decide whether each pair contained the same or different sequence. Participants were then asked whether

they were aware of hearing faint sounds accompanying the flashes. In a final critical task, participants had to detect faint sounds, presented with or without irrelevant visual flashes.

Of the 40 participants, 22 per cent of normal participants reported hearing sounds accompanying the visual flashes in the 'Morse-code' task. These participants performed significantly better in the visual sequence discrimination task, supporting previous suggestions that the ability to recode visual events as sounds may influence visual discrimination of rhythmic sequences. Furthermore, the study is the first to suggest that such auditory-recoding of visual events not only indirectly affects visual performance but directly affect auditory signal detection, making it harder to hear real sounds in the presence of visual stimulation. This kind of visually-evoked auditory response (V-EAR) is thus apparently not only quite common but it can impact on detection of real sounds.

Dr Elliot Freeman, lead author from the Cognitive Neuroscience Research Unit at City, University of London, said:

"Our study has provided the first evidence that a surprisingly high proportion of individuals can sometimes report auditory sensations evoked by visual flashes. This phenomenon is fascinating as it suggests that many of us can use our hearing to help us to remember visual sequences and process them more efficiently. In some people the ability to hear what we see such conversion of images into sounds might compensate for a poorer ability to discriminate pure auditory sequences.

"These internal sounds seem to be perceptually real enough to interfere with the detection of externally-generated sounds. The finding that this 'hearing-motion' phenomenon seems to be much more prevalent compared to other synaesthesias might occur due to the strength of the natural connection between [sound](#) and vision.

More information: Christopher Fassnidge et al. A deafening flash! Visual interference of auditory signal detection, *Consciousness and Cognition* (2017). [DOI: 10.1016/j.concog.2016.12.009](https://doi.org/10.1016/j.concog.2016.12.009)

Take the test here: www.staff.city.ac.uk/~sbbf269/...rough_your_eyes.html

Provided by City University London

Citation: Synaesthesia 'hearing-motion' phenomenon more common than previously thought, says study (2017, January 19) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-01-synaesthesia-hearing-motion-phenomenon-common-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.