

Vision problems after concussion

January 6 2017

Vision problems are a common and sometimes lasting consequence of head injuries—from children and teens with sports-related concussions to military personnel with combat-related traumatic brain injury (TBI). New research and perspectives on TBI and vision are presented in the January feature issue of *Optometry and Vision Science*, official journal of the American Academy of Optometry.

Visual dysfunction after TBI can take many forms and can contribute to lasting disability following [head injuries](#). The feature issue articles add to the growing body of evidence showing important connections between the visual system and brain dysfunction following head injuries. The new research also shows similarities between sports concussions and more severe blast injuries experienced by military service personnel—confirming that an ongoing commitment to TBI research can have far-reaching benefits.

New Insights on Vision Problems after Civilian and Military TBI

Post-concussion visual dysfunction can occur on its own or in association with vestibular dysfunction (balance problems or dizziness). A study led by Mark W. Swanson, OD, MSPH, FAAO, of Children's of Alabama, Birmingham, evaluated the relationship between vision symptoms and [academic problems](#) in children with concussion. The study included 276 children and adolescents with multiple post-concussion symptoms lasting ten days or longer.

A median of three weeks after concussion, 46 percent of the patients had vision symptoms while 29 percent had academic difficulties. Academic problems were twice as likely for children with vision or hearing problems, and 20 times more likely for those with difficulty concentrating.

Among patients who still had post-concussion symptoms after 30 days, the risk of academic problems was three times higher for those with vision symptoms and 15 times higher for those with difficulty concentrating. The authors believe that vision assessment should be part of "return to learn" strategies for children with lasting symptoms after concussion.

A study led by José E. Capó-Aponte, OD, PhD, FAAO of the US Army Aeromedical Research Laboratory assessed the rate of visual dysfunction in 500 military personnel with two types of mild TBI. About two-thirds had blast-related TBI, most often caused by improvised explosive devices. The rest had nonblast-related TBI, with causes such as falls or blunt force.

Nearly one-third of patients had significant visual problems and symptoms one year after injury—whether they had blast-related or nonblast-related TBI. Eye pain was more common in the early stages, while double vision (diplopia) was more frequent later on. The findings highlight the need for ongoing vision rehabilitation care for veterans with TBI. They also suggest that visual problems resulting from civilian TBI—which is rarely blast-related—are similar to those after blast-related military TBI.

In an invited commentary, James F. Jorkasky of the [National Alliance for Eye and Vision Research](#) and vision rehabilitation expert Gregory L. Goodrich, PhD, FAAO, discuss the importance of continued research into [vision](#) problems after head trauma. Since 2009, Congress has

provided annual funding for a dedicated Vision Research Program.

Recognizing the potentially severe impact of concussions—especially repeated concussions—has led to basic research to better understand the visual consequences of TBI, as well as "translational" research to apply those findings to clinical care. The authors write: "Although Vision Program Research immediately addresses deployment-related eye injuries, what it discovers could ultimately apply to civilian eye trauma from accidents and to visual implications from TBIs or sports-related concussive injuries."

More information: [Click here to read the special January issue of Optometry and Vision Science.](#)

Provided by Wolters Kluwer Health

Citation: Vision problems after concussion (2017, January 6) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-01-vision-problems-concussion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--