

Zinc eaten at levels found in biofortified crops reduces 'wear and tear' on DNA

January 3 2017

A new study by researchers from the UCSF Benioff Children's Hospital Research Institute (CHORI) shows that a modest 4 milligrams of extra zinc a day in the diet can have a profound, positive impact on cellular health that helps fight infections and diseases. This amount of zinc is equivalent to what biofortified crops like zinc rice and zinc wheat can add to the diet of vulnerable, nutrient deficient populations.

The study, published in the *American Journal of Clinical Nutrition*, was led by CHORI Senior Scientist Janet King, PhD. King and her team are the first to show that a modest increase in dietary [zinc](#) reduces [oxidative stress](#) and damage to DNA.

"We were pleasantly surprised to see that just a small increase in [dietary zinc](#) can have such a significant impact on how metabolism is carried out throughout the body," says King. "These results present a new strategy for measuring the impact of zinc on health and reinforce the evidence that food-based interventions can improve micronutrient deficiencies worldwide."

Zinc is ubiquitous in our body and facilitates many functions that are essential for preserving life. It plays a vital role in maintaining optimal childhood growth, and in ensuring a healthy immune system. Zinc also helps limit inflammation and oxidative stress in our body, which are associated with the onset of chronic cardiovascular diseases and cancers.

Around much of the world, many households eat polished white rice or

highly refined wheat or maize flours, which provide energy but do not provide enough essential micronutrients such as zinc. Zinc is an essential part of nearly 3,000 different proteins, and it impacts how these proteins regulate every cell in our body. In the absence of sufficient zinc, our ability to repair everyday wear and tear on our DNA is compromised.

In the randomized, controlled, six-week study the scientists measured the impact of zinc on human metabolism by counting DNA strand breaks. They used the parameter of DNA damage to examine the influence of a moderate amount of zinc on healthy living. This was a novel approach, different from the commonly used method of looking at zinc in the blood or using stunting and morbidity for assessing zinc status.

According to King, these results are relevant to the planning and evaluation of food-based solutions for mitigating the impact of hidden hunger and malnutrition. King believes that biofortification can be a sustainable, long-term solution to [zinc deficiency](#).

More information: Sarah J Zyba et al, A moderate increase in dietary zinc reduces DNA strand breaks in leukocytes and alters plasma proteins without changing plasma zinc concentrations, *The American Journal of Clinical Nutrition* (2016). [DOI: 10.3945/ajcn.116.135327](https://doi.org/10.3945/ajcn.116.135327)

Provided by Children's Hospital & Research Center Oakland

Citation: Zinc eaten at levels found in biofortified crops reduces 'wear and tear' on DNA (2017, January 3) retrieved 21 July 2024 from <https://medicalxpress.com/news/2017-01-zinc-eaten-biofortified-crops-dna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.