

# Direct admission to PCI center reduces mortality in STEMI

February 16 2017

---



(HealthDay)—For patients with ST-segment elevation myocardial

infarction (STEMI) treated with percutaneous coronary intervention (PCI), direct admission to a primary PCI center is associated with lower 12-month mortality, according to a study published online Feb. 15 in *JACC: Cardiovascular Interventions*.

Using data from a prospective nationwide registry, Damian Kawecki, M.D., Ph.D., from the Medical University of Silesia in Poland, and colleagues examined the impact of direct admission versus transfer via regional hospital to a PCI center on time delays and 12-month [mortality](#) in STEMI patients. A total of 70,093 patients were eligible for analysis, of whom 56 percent were admitted directly to a PCI center.

The researchers found that direct admission correlated with a 44-minute shorter median symptoms-to-admission time (P

"In a large, community-based cohort of patients with STEMI treated by PCI, direct admission to a primary PCI center was associated with lower 12-month mortality and should be preferred to transfer via a regional non-PCI-capable facility," the authors write.

**More information:** [Full Text](#)  
[Editorial](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Direct admission to PCI center reduces mortality in STEMI (2017, February 16)  
retrieved 26 April 2024 from  
<https://medicalxpress.com/news/2017-02-admission-pci-center-mortality-stemi.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--