

Anterior segment parameters tied to gonioscopic angle closure

February 10 2017



(HealthDay)—Baseline anterior segment parameters are associated with

development of incident gonioscopic angle closure after four years, according to a study published online Feb. 9 in *JAMA Ophthalmology*.

Monisha E. Nongpiur, M.D., Ph.D., from the Singapore Eye Research Institute, and colleagues recruited 342 [participants](#) aged 50 years or older to a prospective observational study. At baseline and after four years, participants underwent gonioscopy and anterior segment optical coherence tomography imaging.

The researchers found that 14.3 percent of participants developed gonioscopic angle closure after four years. These participants had a smaller baseline angle opening distance at 750 μm (AOD750), trabecular iris surface area at 750 μm , anterior chamber area, and anterior chamber volume (all P

"These findings suggest that smaller AOD750 and larger LV [measurements](#) are associated with the development of incident gonioscopic angle closure after four years among participants with gonioscopically open angles at baseline," the authors write.

More information: [Full Text](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Anterior segment parameters tied to gonioscopic angle closure (2017, February 10) retrieved 28 April 2024 from <https://medicalxpress.com/news/2017-02-anterior-segment-parameters-tied-gonioscopic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
