

Children with a communication disorder have poorer sleep, says study

February 7 2017, by George Wigmore



Credit: City University London

Children with a communication disorder such as autism have poorer sleep, compounding the existing language issues that result from such conditions, according to a new study from City, University of London.

The study, which is published in the *International Journal of Developmental Disabilities*, is one of the first to document sleep

behaviours in [children](#) with language impairment, and to look at the relationship between language and sleep.

Professor Nikki Botting, author of the paper and a Professor of Developmental Disorders in the Division of Language and Communication Science at City, University of London, said:

"Sleep patterns relate directly to communication skills and the whole process of sleep is vital for language development and learning. What we found in our study is that impoverished sleep links to poorer language development, even within groups with communication disorders. "

We think that spending more time falling asleep and early waking could potentially interfere with the process of language consolidation especially in the context of an already compromised language system.

Language, communication and sleep

Language and communication are key aspects of cognitive function in everyday life and there is now evidence that sleep plays an important role in consolidating these aspects of development in typical children.

Notably, sleep has been found to boost learning of novel experimental words in childhood and even in babies as young as six months.

Sleep problems are common in children with autism spectrum disorder (ASD) but there is currently little research into [sleep patterns](#) of children with other communication difficulties, such as developmental language disorders.

Children with ASD and Developmental Language Disorders (DLDs) represent a substantial population of children, comprising in 7% and 1% of children respectively. As a result the number of children affected is

significant and it is important that we understand as much as possible about associated difficulties.

To investigate the effects of communication disorders on sleep, the researchers used parental questionnaires to explore the relationship between sleep and language.

There were 65 child participants in total aged between 3-18 years. Parents reported on 28 children with a developmental [communication disorder](#), including eight with ASD and 20 with DLD. There were also 37 questioned who were typical developing.

Questions covered five main areas including demographic information; sleep behaviour; reported [sleep difficulties](#); language and other developmental skills.

The researchers found that the children with a developmental communication disorder showed more [sleep problems](#) than their typical peers particularly in terms of getting to sleep and early waking.

Furthermore, significant correlations were found between sleep behaviour and language for children in both groups.

Professor Nikki Botting said:

"Our study provides evidence for a direct association between everyday sleep behaviours and language ability in a clinical population. "

It does this in two ways, as not only did children with communication disorders show more sleep difficulties than their typically developing peers but sleep behaviours also predicted social-[language](#) difficulties in both groups.

The authors suggest that health professionals involved with families who have communication disorders might consider asking families about sleep patterns, and suggest interventions that involve [sleep](#) behaviour when advising on strategies that might help with [communication](#) difficulties.

More information: Nicola Botting et al. Sleep behaviour relates to language skills in children with and without communication disorders, *International Journal of Developmental Disabilities* (2017). [DOI: 10.1080/20473869.2017.1283766](#)

Provided by City University London

Citation: Children with a communication disorder have poorer sleep, says study (2017, February 7) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-02-children-disorder-poorer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--