


Furosemide plus matched hydration cuts contrast-induced AKI

February 21 2017



(HealthDay)—For high-risk patients undergoing percutaneous coronary

intervention or transcatheter aortic valve replacement, furosemide with matched hydration via the RenalGuard system may reduce the incidence of contrast-induced acute kidney injury (CI-AKI), according to a meta-analysis published in the Feb. 27 issue of *JACC: Cardiovascular Interventions*.

Alessandro Putzu, M.D., from the Cardiocentro Ticino in Lugano, Switzerland, and colleagues conducted a meta-analysis to examine whether furosemide with matched hydration using the RenalGuard System correlated with reduced CI-AKI in patients undergoing interventional procedures. Four trials published from 2011 to 2016 with 698 patients undergoing percutaneous coronary procedures and transcatheter [aortic valve replacement](#) were included in the analysis.

The researchers found that, compared with control treatment, RenalGuard therapy correlated with significantly reduced incidence of CI-AKI (7.76 versus 21.43 percent; odds ratio, 0.31), as well as a reduced need for renal replacement therapy (0.58 versus 3.45 percent; odds ratio, 0.19). There were no major adverse events among patients undergoing RenalGuard therapy.

"Further independent high-quality randomized trials should elucidate the effectiveness and safety of this prophylactic intervention in interventional cardiology," the authors write.

More information: [Abstract](#)
[Full Text](#)
[Editorial](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Furosemide plus matched hydration cuts contrast-induced AKI (2017, February 21)

retrieved 28 April 2024 from

<https://medicalxpress.com/news/2017-02-furosemide-hydration-contrast-induced-aki.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.