

HbA1c, waist-to-height ratio predict dyslipidemia in T1DM

February 1 2017



(HealthDay)—For youth with type 1 diabetes, hemoglobin A1c (A1C)

and waist-to-height ratio (WHtR) are modifiable risk factors that predict change in dyslipidemia, according to a study published online Jan. 26 in *Diabetes Care*.

Amy S. Shah, M.D., from Cincinnati Children's Hospital & University of Cincinnati, and colleagues examined the [risk factors](#) associated with progression and regression of dyslipidemia in 1,478 youths with type 1 diabetes at baseline and at a mean follow-up of 7.1 years.

The researchers found that 19, 5, 69, and 7 percent of youth with type 1 diabetes had non-high-density lipoprotein cholesterol (HDL-C) that progressed, regressed, was stable normal, and stable abnormal, respectively. For HDL-C, the corresponding percentages were 3, 3, 94, and 1 percent. Higher A1C area under the curve (AUC) and higher WHtR AUC in males were associated with non-HDL-C progression. Lower WHtR AUC was associated with non-HDL-regression, while male sex and higher WHtR AUC correlated with HDL-C progression. Due to small numbers, HDL-C regression was not modeled.

"A1C and WHtR are modifiable risk factors associated with change in [dyslipidemia](#) over time in youth with type 1 diabetes," the authors write.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: HbA1c, waist-to-height ratio predict dyslipidemia in T1DM (2017, February 1)
retrieved 25 April 2024 from
<https://medicalxpress.com/news/2017-02-hba1c-waist-to-height-ratio-dyslipidemia-t1dm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.