


The reasons for our left or right-handedness

February 17 2017



Judith Schmitz and Sebastian Ocklenburg are interested in right-left differences.
Credit: Ruhr-Universitaet-Bochum

It is not the brain that determines if people are right or left-handed, but the spinal cord. This has been inferred from the research results compiled by a team headed by private lecturer Dr Sebastian Ocklenburg, Judith Schmitz, and Prof Dr H. C. Onur Güntürkün. Together with

colleagues from the Netherlands and from South Africa, the biopsychologists at Ruhr-Universität Bochum have demonstrated that gene activity in the spinal cord is asymmetrical already in the womb. A preference for the left or the right hand might be traced back to that asymmetry.

"These results fundamentally change our understanding of the cause of hemispheric asymmetries," conclude the authors. The team report about their study in the journal *eLife*.

Preference in the womb

To date, it had been assumed that differences in gene activity of the right and left hemisphere might be responsible for a person's handedness. A preference for moving the left or right hand develops in the womb from the eighth week of pregnancy, according to ultrasound scans carried out in the 1980s. From the 13th week of pregnancy, unborn children prefer to suck either their right or their left thumb.

Arm and hand movements are initiated via the motor cortex in the brain. It sends a corresponding signal to the spinal cord, which in turn translates the command into a motion. The [motor cortex](#), however, is not connected to the spinal cord from the beginning. Even before the connection forms, precursors of handedness become apparent. This is why the researchers have assumed that the cause of right respective left preference must be rooted in the spinal cord rather than in the brain.

The influence of environmental factors

The researchers analysed the gene expression in the spinal cord during the eighth to twelfth week of pregnancy and detected marked right-left differences in the eighth week – in precisely those spinal cord segments

that control the movements of arms and legs. Another study had shown that unborn children carry out asymmetric [hand movements](#) just as early as that.

The researchers, moreover, traced the cause of asymmetric [gene activity](#). Epigenetic factors appear to be at the root of it, reflecting environmental influences. Those influences might, for example, lead to enzymes bonding methyl groups to the DNA, which in turn would affect and minimise the reading of genes. As this occurs to a different extent in the left and the right [spinal cord](#), there is a difference to the activity of genes on both sides.

More information: Sebastian Ocklenburg et al. Epigenetic regulation of lateralized fetal spinal gene expression underlies hemispheric asymmetries, *eLife* (2017). [DOI: 10.7554/eLife.22784](https://doi.org/10.7554/eLife.22784)

Provided by Ruhr-Universitaet-Bochum

Citation: The reasons for our left or right-handedness (2017, February 17) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-02-left-right-handedness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
